

DECOUPLING OF MIXED METHODS BASED ON GENERALIZED HELMHOLTZ DECOMPOSITIONS*

LONG CHEN[†] AND XUEHAI HUANG[‡]

Abstract. A framework to systematically decouple high order elliptic equations into a combination of Poisson-type and Stokes-type equations is developed. The key is to systematically construct the underlying commutative diagrams involving the complexes and Helmholtz decompositions in a general way. Discretizing the decoupled formulation leads to a natural superconvergence between the Galerkin projection and the decoupled approximation. Examples include but are not limited to the primal formulations and mixed formulations of the biharmonic equation, fourth order curl equation, and triharmonic equation. As a byproduct, Helmholtz decompositions for many dual spaces are obtained.

Key words. differential complex, commutative diagram, Helmholtz decomposition, mixed formulation, decoupling, discretization

AMS subject classifications. 58J10, 65N12, 65N22, 65N30

DOI. 10.1137/17M1145872

1. Introduction. We shall develop a framework to systematically decouple high order elliptic equations into a combination of Poisson-type and Stokes-type equations. The key is to systematically construct the underlying commutative diagrams involving the complexes and Helmholtz decompositions in a general way.

Differential complexes and corresponding Helmholtz decompositions play a fundamental role in the design and analysis of mixed finite element methods. Among many others, the de Rham complex for the Hodge Laplacian and the elasticity complex for the linear elasticity are two successful examples [4, 5]. A direct and useful result of a differential complex is the Helmholtz decomposition. With this decomposition, the kernel spaces of differential operators involved in the complex are characterized clearly. The generalized Helmholtz decomposition of Banach spaces presented in this paper can be regarded as a generalization of the well-known Helmholtz–Hodge decomposition in [4]. The Helmholtz–Hodge decomposition is L^2 -orthogonal, while the generalized Helmholtz decomposition is a direct sum but not necessarily orthogonal.

*Received by the editors September 1, 2017; accepted for publication (in revised form) June 29, 2018; published electronically September 11, 2018.

<http://www.siam.org/journals/sinum/56-5/M114587.html>

Funding: The first author was supported by NSF grant DMS-1418934 and in part by the Sea Poly Project of Beijing Overseas Talents. The second author was supported by the NSFC Projects 11771338 and 11671304, Zhejiang Provincial Natural Science Foundation of China Projects LY17A010010 and LY15A010015, and Wenzhou Science and Technology Plan Project G20160019.

[†]Beijing Institute for Scientific and Engineering Computing, Beijing University of Technology, Beijing 100124, China, and Department of Mathematics, University of California at Irvine, Irvine, CA 92697 (chenlong@math.uci.edu).

[‡]Corresponding author. School of Mathematics, Shanghai University of Finance and Economics, Shanghai 200433, China (xuehaihuang@gmail.com).

Our approach is based on the following diagram:

$$(1.1) \quad \begin{array}{ccccccc} & & X & \xrightarrow{J_X} & X' & & \\ & & & & & \cup & \\ & & & & & \Sigma & \\ 0 & \longrightarrow & P & \xrightarrow{d^-} & \Sigma & \xrightarrow{d} & V' \longrightarrow 0 \\ & & & & \Pi_\Sigma \uparrow & & \uparrow J_V \\ & & & & \tilde{\Sigma} & \xleftarrow{\Pi_V} & V \end{array} ,$$

where the isomorphisms J_X and J_V are the inverses of the Riesz representations, and the rest of the linear operators are all continuous but not necessarily isomorphic. The middle complex is exact, i.e., $\ker(d) = \text{img}(d^-)$, d is surjective, and d^- is injective. A stable Helmholtz decomposition can be derived from (1.1):

$$(1.2) \quad \Sigma = d^- P \oplus \Pi_\Sigma \Pi_V V.$$

By finding the specific diagrams of (1.1), we recover several well-known Helmholtz decompositions and discover many new Helmholtz decompositions. In particular, we obtain Helmholtz decompositions for many dual spaces. We summarize the details of these Helmholtz decompositions in Table 1.

TABLE 1
Examples of Helmholtz decomposition generated from (1.1)–(1.2).

Hilbert space	Helmholtz decomposition	Refs.
$L^2(\Omega)$ in 2D and 3D	$\nabla H_0^1(\Omega) \oplus^\perp \text{curl } \mathbf{H}^1(\Omega)$	[4, 29, 18]
$\mathbf{H}^{-1}(\Omega)$ in 2D	$\nabla L_0^2(\Omega) \oplus^\perp \Delta(\text{curl } H_0^2(\Omega))$	[45]
$\mathbf{H}^{-1}(\text{div}, \Omega)$ in 2D	$\nabla H_0^1(\Omega) \oplus \text{curl } L^2(\Omega)$	[14]
$\mathbf{H}^{-1}(\text{div div}, \Omega; \mathbb{S})$ in 2D	$\nabla^s \times \mathbf{H}^1(\Omega; \mathbb{R}^2) \oplus \pi H_0^1(\Omega)$	[40]
$\mathbf{H}(\text{div div}, \Omega; \mathbb{S})$ in 2D	$\nabla^s \times \mathbf{H}^1(\Omega; \mathbb{R}^2) \oplus \pi \Delta^{-1} L^2(\Omega)$	section 2.4
$\mathbf{H}^{-2}(\text{rot rot}, \Omega; \mathbb{S})$ in 2D	$\boldsymbol{\varepsilon} L^2(\Omega; \mathbb{R}^2) \oplus \text{curl curl } H_0^2(\Omega)$	section 2.5
$\mathbf{H}^{-2}(\text{div}^3, \Omega)$ in 2D	$\text{sym curl } \mathbf{H}^1(\Omega; \mathbb{S}) \oplus \Xi \nabla H_0^2(\Omega)$	section 2.7
$\mathbf{H}^{-1}(\text{div}, \Omega)$ in 3D	$\nabla H_0^1(\Omega) \oplus \text{curl } L^2(\Omega; \mathbb{R}^3)$	section 2.2
$\mathbf{H}^{-1}(\text{curl}, \Omega)$ in 3D	$\nabla L_0^2(\Omega) \oplus \text{curl } \mathbf{H}_0(\text{curl}, \Omega)$	section 2.3
$\mathbf{H}(\text{curl curl}, (K_0^c)')$ in 3D	$\text{spn}^{-1} \text{skw } \mathbf{H}^0(\text{div}, \text{dev sym}) \oplus K_0^c$	section 2.6

An abstract two-term mixed formulation based on the commutative diagram (1.1) is as follows: given $g \in \Sigma'$ and $f \in V'$, find $(\sigma, u) \in \Sigma \times V$ such that

$$(1.3) \quad (\sigma, \tau)_{X'} + \langle d\tau, u \rangle = \langle g, \tau \rangle \quad \forall \tau \in \Sigma,$$

$$(1.4) \quad \langle d\sigma, v \rangle = \langle f, v \rangle \quad \forall v \in V.$$

Under the assumption that the norm equivalence

$$(1.5) \quad \|\tau\|_\Sigma^2 \approx \|\tau\|_{X'}^2 + \|d\tau\|_{V'}^2, \quad \forall \tau \in \Sigma$$

holds, the mixed formulation (1.3)–(1.4) is well-posed. Indeed, d is surjective from the commutative diagram (1.1), and the norm equivalence (1.5) can guarantee the continuity of the bilinear forms and the coercivity of $(\cdot, \cdot)_{X'}$ on $\ker(d)$.

To discretize the inner product $(\sigma, \tau)_{X'}$, we introduce $\phi = J_X^{-1} \sigma \in X$ and obtain an equivalent but unfolded three-term formulation: find $(\phi, u, \sigma) \in X \times V \times \Sigma$ such

that

$$(1.6) \quad (\phi, \psi)_X - \langle \sigma, d'v + \psi \rangle = -\langle f, v \rangle \quad \forall (\psi, v) \in X \times V,$$

$$(1.7) \quad \langle d'u + \phi, \tau \rangle = \langle g, \tau \rangle \quad \forall \tau \in \Sigma.$$

Equation (1.6) is a combination of (1.4) and $\phi = J_X^{-1}\sigma$, and (1.7) follows from (1.3) and $\phi = J_X^{-1}\sigma$.

Applying the Helmholtz decomposition (1.2) to the unfolded formulation (1.6)–(1.7), we obtain a decoupled formulation: find $w, u \in V$, $\phi \in X$, and $p \in P/\ker d^-$ such that

$$(1.8) \quad (w, v)_V = \langle f, v \rangle \quad \forall v \in V,$$

$$(1.9) \quad (\phi, \psi)_X - \langle d^-p, \psi \rangle = \langle \Pi_\Sigma \Pi_V w, \psi \rangle \quad \forall \psi \in X,$$

$$(1.10) \quad \langle d^-q, \phi \rangle = \langle g, d^-q \rangle \quad \forall q \in P/\ker d^-,$$

$$(1.11) \quad (u, \chi)_V = \langle g - \phi, \Pi_\Sigma \Pi_V \chi \rangle \quad \forall \chi \in V.$$

The middle system (1.9)–(1.10) of (ϕ, p) is now a Stokes-type system, and (1.8) and (1.11) are usually Poisson-type equations depending on the inner product $(\cdot, \cdot)_V$.

By finding the underlying complexes, we recover some recent results on the decoupling of

- the HHJ method for the plate problem into two Poisson equations and one linear elasticity problem [40];
- the primal formulation of the biharmonic equation in two and three dimensions into two Poisson equations and one Stokes-type equation [36, 38, 27];
- the primal formulation of the fourth order elliptic singular perturbation problem into two Poisson equations and one Brinkman problem [27];
- the primal formulation of the fourth order curl equation into two Maxwell equations and one Stokes equation [52].

Moreover, we can get new decouplings by using the framework developed in this paper; for example, we decouple

- the mixed formulation of the fourth order curl equation into two Maxwell equations and one mixed formulation of the Poisson-type equation in section 3.5;
- the primal formulation of the triharmonic equation in two dimensions into two biharmonic equations and one tensorial Stokes equation in section 3.6;
- the mixed formulation of the triharmonic equation in two dimensions into two biharmonic equations and one tensorial Poisson-type equation in section 3.7;
- the m th harmonic equation into two $(m - 1)$ th harmonic equations and one tensorial Stokes-type equation in section 3.6.

Compared to the original formulation, it is much easier to construct conforming finite element spaces for the decoupled formulation, since the order of the system is reduced and the finite element methods for the Stokes equation and the Poisson equation are well developed. We shall also show a natural superconvergence between the Galerkin projection and the approximation based on the decoupled formulation.

We are motivated by the pioneering work of such decoupling for the Reissner–Mindlin plate model [14], which has existed since the 1980s, and a recent decomposition for the HHJ formulation of the Kirchhoff plate model [40, 47] and the biharmonic equation in three dimensions [46]. Results can be also found for the primal formulation of biharmonic equations [36, 38, 27], the primal formulation of the fourth order curl equation [52, 11], the eigenvalue problem of biharmonic equation [54], and the

linear second order elliptic problem in nondivergence form [28]. Viewed as the polyharmonic generalized Stokes problem, the m th harmonic equation was decoupled into $(2m - 2)$ Poisson-type problems and one generalized Stokes equation over the symmetric tensors by applying a split recursively in [27]. As comparison, we decouple the m th harmonic equation into two $(m - 1)$ th harmonic equations and one tensorial Stokes-type equation independently, and recursively decouple into 2^{m-1} Poisson equations and $(2^{m-1} - 1)$ Stokes-type equations, which is different from the decoupling in [27] for $m \geq 3$. We can also stop the decoupling at biharmonic equations, which can be discretized directly by many existing finite elements methods. We refer the reader to [48, 53] for more on reducing the m th harmonic equation into lower order partial differential equations. Our framework unifies most of those results and will lead to many more Hodge decompositions, especially for high order elliptic equations. This way, we can decouple the higher order partial differential equations into lower order ones, which makes the discretization easier.

In addition, the explicit expression of the kernel space can be used to develop fast solvers; see, for example, [33, 3, 35, 20, 21]. The Helmholtz decomposition is also a key tool to construct the a posteriori error estimators of nonconforming and mixed finite element methods [1, 16, 17, 19, 37, 22]. The important role of the structure revealed in our work for designing fast solvers and the a posterior error analysis will be explored somewhere else.

The rest of this paper is organized as follows. In section 2, we establish the generalized Helmholtz decomposition based on the commutative diagram and give several examples. The abstract mixed formulation and its decomposition based on the Helmholtz decomposition are presented in section 3. In section 4, we discretize the decoupled formulation directly illustrated by two examples. Throughout this paper, we use “ $\lesssim \dots$ ” to mean that “ $\leq C \dots$ ”, where C is a generic positive constant independent of mesh-size h , which may take different values at different appearances. And $a \approx b$ means $a \lesssim b$ and $b \lesssim a$.

2. Generalized Helmholtz decompositions. In this section we apply the splitting lemma of Banach spaces and Hilbert spaces to differential complexes and obtain Helmholtz-type decompositions for several Sobolev spaces with negative index.

2.1. Background in functional analysis. We start from a short exact sequence

$$(2.1) \quad \widetilde{W} \xrightarrow{\tilde{d}_2} \widetilde{V} \xrightarrow{\tilde{d}_1} \widetilde{U} \longrightarrow 0.$$

Here the capital letters represent Banach spaces and \tilde{d}_i ($i = 1, 2$) are bounded linear operators. The sequence (2.1) is exact, meaning that

$$\ker(\tilde{d}_1) = \text{img}(\tilde{d}_2), \quad \text{img}(\tilde{d}_1) = \widetilde{U}.$$

The space \widetilde{W} can be further reduced to the quotient space $\widetilde{W} / \ker(\tilde{d}_2)$ so that

$$0 \longrightarrow \widetilde{W} / \ker(\tilde{d}_2) \xrightarrow{\tilde{d}_2} \widetilde{V} \xrightarrow{\tilde{d}_1} \widetilde{U} \longrightarrow 0$$

forms a short exact sequence in the context of group [30].

Let U, V be two additional Banach spaces and $d_1 : U \rightarrow V$ be a bounded linear operator. Let $I_V : d_1U \rightarrow \tilde{V}$ be a bounded linear operator and $J_U : U \rightarrow \tilde{U}$ be an isomorphism satisfying the assumption

$$(2.2) \quad \tilde{d}_1 I_V d_1 u = J_U u \quad \forall u \in U,$$

which can be summarized as the following commutative diagram:

$$(2.3) \quad \begin{array}{ccccccc} \tilde{W} & \xrightarrow{\tilde{d}_2} & \tilde{V} & \xrightarrow{\tilde{d}_1} & \tilde{U} & \longrightarrow & 0 \\ & & \uparrow I_V & & \uparrow J_U & & \\ & & V & \xleftarrow{d_1} & U & & \end{array} .$$

To derive the generalized Helmholtz decomposition, we first recall the splitting lemma in algebraic topology.

LEMMA 2.1 (splitting lemma in [30]). *For a short exact sequence*

$$0 \longrightarrow U \xrightarrow{d_1} V \xrightarrow{d_2} W \longrightarrow 0$$

of abelian groups the following statements are equivalent:

- (a) *There is a homomorphism $d_3 : V \rightarrow U$ such that $d_3 d_1$ is the identity on U .*
- (b) *There is a homomorphism $d_4 : W \rightarrow V$ such that $d_2 d_4$ is the identity on W .*
- (c) *The group V is isomorphic to the direct sum of U and W , with d_1 corresponding to the natural injection of U and d_2 to the natural projection onto W .*

Apparently Banach spaces are abelian groups under addition. Then we have a generalized Helmholtz decomposition as follows.

THEOREM 2.2. *Suppose we have a short exact sequence (2.1). Assume the commutative diagram (2.3) holds with all the linear operators being bounded and $J_U : U \rightarrow \tilde{U}$ being an isomorphism. Then we have a stable Helmholtz decomposition,*

$$(2.4) \quad \tilde{V} = \tilde{d}_2(\tilde{W}/\ker(\tilde{d}_2)) \oplus I_V d_1 U,$$

where \oplus means the direct sum. More precisely, for any $\tilde{v} \in \tilde{V}$, there exist $\tilde{w} \in \tilde{W}/\ker \tilde{d}_2$ and $u \in U$ such that

$$(2.5) \quad \tilde{v} = \tilde{d}_2 \tilde{w} + I_V d_1 u,$$

$$(2.6) \quad \|\tilde{w}\|_{\tilde{W}} + \|u\|_U \lesssim \|\tilde{v}\|_{\tilde{V}}.$$

Proof. By assumptions, we have a short exact sequence

$$0 \longrightarrow \tilde{W}/\ker(\tilde{d}_2) \xrightarrow{\tilde{d}_2} \tilde{V} \xrightarrow{J_U^{-1} \tilde{d}_1} U \longrightarrow 0.$$

And $I_V d_1$ forms a right inverse of $J_U^{-1} \tilde{d}_1$. Thus (2.4) holds by Lemma 2.1, which is stable as all operators involved are continuous. \square

The generalized Helmholtz decomposition (2.4) is a direct result of the splitting lemma after finding the underlying commutative diagram (2.3). One contribution of this paper is to construct various commutative diagrams to induce stable Helmholtz decompositions on Sobolev spaces of negative order.

Remark 2.3. Consider Hilbert space \tilde{V} with inner product $(\cdot, \cdot)_{\tilde{V}}$. The quotient space $\ker(\tilde{d}_1)/\tilde{d}_2(\tilde{W})$ is isomorphic to the space of harmonic forms

$$\mathfrak{H} := \left\{ \tilde{v} \in \tilde{V} : \tilde{d}_1\tilde{v} = 0, (\tilde{v}, \tilde{d}_2\tilde{w})_{\tilde{V}} = 0 \quad \forall \tilde{w} \in \tilde{W} \right\}.$$

When the quotient space $\ker(\tilde{d}_1)/\tilde{d}_2(\tilde{W})$ is nontrivial, i.e., the sequence (2.1) is not exact, the Helmholtz decomposition will be

$$\tilde{V} = \tilde{d}_2(\tilde{W}/\ker(\tilde{d}_2)) \oplus I_V d_1U \oplus \mathfrak{H}.$$

In Theorem 2.2, the decomposition is a direct sum but not necessarily orthogonal. Indeed, in the proof we do not use the inner product structure. We now explore the orthogonality for Hilbert complexes. In what follows, we always denote by $\langle \cdot, \cdot \rangle$ the duality pairing and reserve (\cdot, \cdot) for the L^2 inner product.

Denote by X' the dual space of a linear space X and by $T' : Y' \rightarrow X'$ the dual of a linear operator $T : X \rightarrow Y$ defined as

$$\langle T'g, x \rangle := \langle g, Tx \rangle.$$

When X is a Hilbert space with an inner product $(\cdot, \cdot)_X$ and X' is the continuous dual of X , by the Riesz representation theorem, we have an isomorphism $J_X : X \rightarrow X'$: for any $w \in X$ and define $J_X w \in X'$ as

$$(2.7) \quad \langle J_X w, v \rangle = (w, v)_X \quad \forall v \in X.$$

The induced inner product and norm for any $w', v' \in X'$ are given by

$$(2.8) \quad \begin{aligned} (w', v')_{X'} &:= (J_X^{-1}w', J_X^{-1}v')_X = \langle J_X^{-1}w', v' \rangle = \langle w', J_X^{-1}v' \rangle, \\ \|w'\|_{X'} &:= \|J_X^{-1}w'\|_X. \end{aligned}$$

Let U, V, W be Hilbert spaces. Suppose we have a short exact sequence of their dual spaces

$$(2.9) \quad 0 \longrightarrow W'/\ker(d'_2) \xrightarrow{d'_2} V' \xrightarrow{d'_1} U' \longrightarrow 0.$$

By Remark 2.15 in [46], the dual complex (2.9) implies the exact sequence

$$(2.10) \quad 0 \longrightarrow U \xrightarrow{d_1} V \xrightarrow{d_2} W.$$

Then it is apparent that $d'_1 J_V d_1$ is an isomorphism from U to U' . By taking $J_U = d'_1 J_V d_1$, the assumption (2.2) holds. With \tilde{X} ($X = U, V, W$) and \tilde{d}_i ($i = 1, 2$) replaced by X' ($X = U, V, W$) and d'_i ($i = 1, 2$), the commutative diagram (2.3) becomes

$$(2.11) \quad \begin{array}{ccccc} W' & \xrightarrow{d'_2} & V' & \xrightarrow{d'_1} & U' & \longrightarrow & 0 \\ & & \uparrow J_V & & \uparrow J_U & & \\ & & V & \xleftarrow{d_1} & U & & \end{array} .$$

Applying Theorem 2.2 to the commutative diagram (2.11), we recover the following orthogonal Helmholtz decomposition.

COROLLARY 2.4. *Suppose the short exact Hilbert sequence (2.9) holds. Then we have the $(\cdot, \cdot)_{V'}$ -orthogonal Helmholtz decomposition*

$$(2.12) \quad V' = d'_2(W'/\ker(d'_2)) \oplus^\perp J_V d_1U.$$

That is, for any $v' \in V'$, there exist $w' \in W'/\ker d'_2$ and $u \in U$ such that

$$(2.13) \quad v' = d'_2w' + J_V d_1u,$$

$$(2.14) \quad \|v'\|_{V'}^2 = \|d'_2w'\|_{V'}^2 + \|J_V d_1u\|_{V'}^2.$$

Corollary 2.4 is indeed one way to express the well-known closed range theorem [51] for Hilbert spaces and the associated orthogonal space decomposition.

In the decomposition (2.12), we need to know J_V , which involves the inner product of Hilbert space V . Sometimes we do not exactly know the space V or do not necessarily need to know. We give an example to illustrate this point.

Consider a dense subspace V of a larger space Y endowed with the inner product $(\cdot, \cdot)_Y$. In most places in this paper, Y is the L^2 space for scalar or vector functions with $(\cdot, \cdot)_Y = (\cdot, \cdot)$ being the L^2 -inner product. We can equip V with the graph inner product

$$(2.15) \quad (w, v)_V := (w, v)_Y + (d_2w, d_2v)_W.$$

Or we can start from Y and define V as the subspace of Y with $\|\cdot\|_V < \infty$. By identifying Y' with Y using the inner product $(\cdot, \cdot)_Y$, we have the rigged Hilbert space [12, 26]

$$(2.16) \quad V \subset Y \subset V'.$$

We compute $J_V : V \rightarrow V'$ as follows: for any $u \in U$ and $v \in V$

$$\langle J_V d_1u, v \rangle = (d_1u, v)_V = (d_1u, v)_Y + (d_2d_1u, d_2v)_W = (d_1u, v)_Y.$$

Thus J_V is the composition of the natural inclusions in (2.16) on d_1U . On the other hand, we can use the commutative diagram to characterize the dual space.

COROLLARY 2.5. *Suppose the short exact Hilbert sequence (2.9) holds, and V is equipped with the graph inner product (2.15). Assume we have another short exact Hilbert sequence*

$$W' \xrightarrow{d'_2} \tilde{V} \xrightarrow{d'_1} U' \longrightarrow 0$$

and a commutative diagram

$$\begin{array}{ccccc} W' & \xrightarrow{d'_2} & \tilde{V} & \xrightarrow{d'_1} & U' \longrightarrow 0 \\ & & \uparrow I & & \uparrow J_U \\ & & V & \xleftarrow{d_1} & U \end{array}$$

with I being the embedding operator. Then $\tilde{V} = V' = d'_2(W'/\ker(d'_2)) \oplus d_1U$.

Proof. The result immediately follows from the previous illustration, Corollary 2.4, and Theorem 2.2. □

We shall present examples in what follows. Let $\Omega \subset \mathbb{R}^n, n = 2, 3$, be a bounded Lipschitz domain. Denote by \mathbb{M} the space of all $n \times n$ tensors, by \mathbb{S} the space of all symmetric $n \times n$ tensors, and by \mathbb{K} the space of all skew-symmetric $n \times n$ tensors. For any tensor $\boldsymbol{\tau} \in \mathbb{M}$, let $\text{sym } \boldsymbol{\tau} := (\boldsymbol{\tau} + \boldsymbol{\tau}^\top)/2$ be the symmetric part of the tensor, and let $\text{skw } \boldsymbol{\tau} := (\boldsymbol{\tau} - \boldsymbol{\tau}^\top)/2$ be the skew-symmetric part. Denote the deviatoric part and the trace of the tensor $\boldsymbol{\tau}$ by $\text{dev } \boldsymbol{\tau}$ and $\text{tr } \boldsymbol{\tau}$ accordingly. We have

$$\text{dev } \boldsymbol{\tau} = \boldsymbol{\tau} - \frac{1}{n}(\text{tr } \boldsymbol{\tau})\mathbf{I}.$$

Define operator $\text{spn} : \mathbb{R}^3 \rightarrow \mathbb{M}$ as follows: for any vector $\mathbf{a} = (a_1, a_2, a_3)^\top \in \mathbb{R}^3$, the tensor $\text{spn } \mathbf{a} \in \mathbb{M}$ is given by

$$\text{spn } \mathbf{a} := \begin{pmatrix} 0 & -a_3 & a_2 \\ a_3 & 0 & -a_1 \\ -a_2 & a_1 & 0 \end{pmatrix}.$$

Denote by \mathfrak{S}_3 the set of all permutations of $(1, 2, 3)$. Define the set of symmetric third order tensors as (cf. [48, section 2])

$$\mathbb{S}(3) := \{ \boldsymbol{\tau} \in (\mathbb{R}^2)^3 : \boldsymbol{\tau}_{j_1, j_2, j_3} = \boldsymbol{\tau}_{j_{\sigma(1)}, j_{\sigma(2)}, j_{\sigma(3)}} \quad \forall (j_1, j_2, j_3) \in \{1, 2\}^3, \sigma \in \mathfrak{S}_3 \}.$$

The symmetric part $\text{sym } \boldsymbol{\tau} \in \mathbb{S}(3)$ of a tensor $\boldsymbol{\tau} \in (\mathbb{R}^2)^3$ is defined by

$$(\text{sym } \boldsymbol{\tau})_{j_1, j_2, j_3} := \frac{1}{\#(\mathbb{S}(3))} \sum_{\sigma \in \mathbb{S}(3)} \boldsymbol{\tau}_{j_{\sigma(1)}, j_{\sigma(2)}, j_{\sigma(3)}}$$

for all $(j_1, j_2, j_3) \in \{1, 2\}^3$. We use standard notation for Sobolev spaces and boldface letters for vector and tensor valued spaces. When we want to emphasize the spatial dimension, we include \mathbb{R}^n in the notation of spaces.

Recall the de Rham complexes in two dimensions,

$$(2.17) \quad 0 \longrightarrow H_0^1(\Omega) \xrightarrow{\text{curl}} \mathbf{H}_0(\text{div}, \Omega) \xrightarrow{\text{div}} L_0^2(\Omega) \longrightarrow 0,$$

$$(2.18) \quad \mathbb{R} \longrightarrow H^1(\Omega) \xrightarrow{\text{curl}} \mathbf{H}(\text{div}, \Omega) \xrightarrow{\text{div}} L^2(\Omega) \longrightarrow 0,$$

$$(2.19) \quad 0 \longrightarrow H_0^{s+2}(\Omega) \xrightarrow{\text{curl}} \mathbf{H}_0^{s+1}(\Omega; \mathbb{R}^2) \xrightarrow{\text{div}} H_0^s(\Omega) \longrightarrow 0,$$

$$(2.20) \quad \mathbb{R} \longrightarrow H^{s+2}(\Omega) \xrightarrow{\text{curl}} \mathbf{H}^{s+1}(\Omega; \mathbb{R}^2) \xrightarrow{\text{div}} H^s(\Omega) \longrightarrow 0,$$

and the de Rham complexes in three dimensions,

$$(2.21) \quad 0 \longrightarrow H_0^1(\Omega) \xrightarrow{\text{grad}} \mathbf{H}_0(\text{curl}, \Omega) \xrightarrow{\text{curl}} \mathbf{H}_0(\text{div}, \Omega) \xrightarrow{\text{div}} L_0^2(\Omega) \longrightarrow 0,$$

$$(2.22) \quad \mathbb{R} \longrightarrow H^1(\Omega) \xrightarrow{\text{grad}} \mathbf{H}(\text{curl}, \Omega) \xrightarrow{\text{curl}} \mathbf{H}(\text{div}, \Omega) \xrightarrow{\text{div}} L^2(\Omega) \longrightarrow 0,$$

$$(2.23) \quad 0 \longrightarrow H_0^{s+3}(\Omega) \xrightarrow{\text{grad}} \mathbf{H}_0^{s+2}(\Omega; \mathbb{R}^3) \xrightarrow{\text{curl}} \mathbf{H}_0^{s+1}(\Omega; \mathbb{R}^3) \xrightarrow{\text{div}} H_0^s(\Omega) \longrightarrow 0,$$

$$(2.24) \quad \mathbb{R} \longrightarrow H^{s+3}(\Omega) \xrightarrow{\text{grad}} \mathbf{H}^{s+2}(\Omega; \mathbb{R}^3) \xrightarrow{\text{curl}} \mathbf{H}^{s+1}(\Omega; \mathbb{R}^3) \xrightarrow{\text{div}} H^s(\Omega) \longrightarrow 0,$$

with $s \in \mathbb{R}$. In 2D, as curl is a rotation of grad operator, we could have similar sequences by replacing curl by grad, div by rot, and $\mathbf{H}(\text{div}, \Omega)$ space by $\mathbf{H}(\text{rot}, \Omega)$ space. For example, an analogue of (2.20) is

$$(2.25) \quad \mathbb{R} \longrightarrow H^{s+2}(\Omega) \xrightarrow{\text{grad}} \mathbf{H}^{s+1}(\Omega; \mathbb{R}^2) \xrightarrow{\text{rot}} H^s(\Omega) \longrightarrow 0.$$

When Ω is simply connected with connected boundary, the L^2 de Rham complexes (2.17)–(2.18) and (2.21)–(2.22) are exact [29, 4, 5], the complexes (2.20) and (2.24)–(2.25) are exact if s is an integer, and the complexes (2.19) and (2.23) are exact if s is a nonnegative integer [46, 25]. When Ω is a bounded domain star-like with respect to a ball, the complexes (2.20) and (2.24)–(2.25) are exact for any $s \in \mathbb{R}$, and the complexes (2.19) and (2.23) are exact if s is nonnegative and $s - \frac{1}{2}$ is not an integer [25, p. 301]. Hereafter, we only consider the domain Ω without harmonic forms; i.e., we always assume the bounded Lipschitz domain Ω is simply connected with the connected boundary in this paper. It will be more sophisticated in the discretization of decoupled formulations and in designing fast solvers and adaptive algorithms when the nontrivial harmonic forms appear in the Helmholtz decompositions. For ease of presentation, we use $\mathbf{H}^s(\Omega)$ to denote $\mathbf{H}^s(\Omega; \mathbb{R}^n)$ for $n = 1, 2, 3$.

We recall the well-known L^2 -orthogonal Helmholtz decomposition (cf. [49, 4])

$$\mathbf{L}^2(\Omega) = \nabla H_0^1(\Omega) \oplus^\perp \text{curl}(\mathbf{H}^1(\Omega)/\ker(\text{curl}))$$

in two and three dimensions from the exact sequences (2.20) and (2.24) with $s = -1$, and the H^{-1} -orthogonal decomposition of $\mathbf{H}^{-1}(\Omega; \mathbb{R}^2)$ (cf. [45, Lemma 2.4])

$$\mathbf{H}^{-1}(\Omega; \mathbb{R}^2) = \nabla L_0^2(\Omega) \oplus^\perp \Delta(\text{curl} H_0^2(\Omega))$$

from the exact sequences (2.25) with $s = -2$. In the following, we present several less well-known and some new Helmholtz decompositions.

2.2. Helmholtz decomposition of $\mathbf{H}^{-1}(\text{div})$ space. For $n = 2$ and 3, define

$$\mathbf{H}^{-1}(\text{div}, \Omega) := \{\phi \in \mathbf{H}^{-1}(\Omega) : \text{div} \phi \in H^{-1}(\Omega)\}$$

with squared norm $\|\phi\|_{\mathbf{H}^{-1}(\text{div})}^2 := \|\phi\|_{-1}^2 + \|\text{div} \phi\|_{-1}^2$.

LEMMA 2.6. *The complex*

$$\mathbf{L}^2(\Omega) \xrightarrow{\text{curl}} \mathbf{H}^{-1}(\text{div}, \Omega) \xrightarrow{\text{div}} H^{-1}(\Omega) \longrightarrow 0$$

is exact in both two and three dimensions.

Proof. We know $\ker(\text{div}) \cap \mathbf{H}^{-1}(\Omega) = \text{curl}(\mathbf{L}^2(\Omega))$ from the exact sequence (2.20) or (2.24) with $s = -2$. Obviously $\ker(\text{div}) \cap \mathbf{H}^{-1}(\text{div}, \Omega) = \ker(\text{div}) \cap \mathbf{H}^{-1}(\Omega)$.

With $s = -1$, it holds that $\text{div} \mathbf{L}^2(\Omega; \mathbb{R}^n) = H^{-1}(\Omega)$, which together with $\mathbf{L}^2(\Omega; \mathbb{R}^n) \subset \mathbf{H}^{-1}(\text{div}, \Omega)$ indicates that $\text{div} \mathbf{H}^{-1}(\text{div}, \Omega) = H^{-1}(\Omega)$. \square

With this exact sequence, we build up the commutative diagram

$$(2.26) \quad \begin{array}{ccccc} \mathbf{L}^2(\Omega) & \xrightarrow{\text{curl}} & \mathbf{H}^{-1}(\text{div}, \Omega) & \xrightarrow{\text{div}} & H^{-1}(\Omega) & \longrightarrow & 0 \\ & & \uparrow I & & \uparrow \Delta & & \\ & & \mathbf{H}_0(\text{curl}, \Omega) & \xleftarrow{\text{grad}} & H_0^1(\Omega) & & \end{array} .$$

By Theorem 2.2, we obtain the Helmholtz decomposition in both two and three dimensions,

$$(2.27) \quad \mathbf{H}^{-1}(\operatorname{div}, \Omega) = \nabla H_0^1(\Omega) \oplus \operatorname{curl}(\mathbf{L}^2(\Omega) / \ker(\operatorname{curl})).$$

In two dimensions, $\ker(\operatorname{curl}) = \mathbb{R}$ and thus the Helmholtz decomposition (2.27) reads as

$$\mathbf{H}^{-1}(\operatorname{div}, \Omega) = \nabla H_0^1(\Omega) \oplus \operatorname{curl} L_0^2(\Omega),$$

which has been presented in [14, Proposition 2.3]. In three dimensions, $\ker(\operatorname{curl}) = \nabla H^1(\Omega)$, it becomes

$$\mathbf{H}^{-1}(\operatorname{div}, \Omega) = \nabla H_0^1(\Omega) \oplus \operatorname{curl}(\mathbf{L}^2(\Omega; \mathbb{R}^3) / \nabla H^1(\Omega)).$$

Applying Remark 2.15 in [46] to the short exact sequence

$$0 \longrightarrow H_0^1(\Omega) \xrightarrow{\operatorname{grad}} \mathbf{H}_0(\operatorname{curl}, \Omega) \xrightarrow{\operatorname{curl}} \mathbf{L}^2(\Omega),$$

we know the dual complex

$$\mathbf{L}^2(\Omega) \xrightarrow{\operatorname{curl}} (\mathbf{H}_0(\operatorname{curl}, \Omega))' \xrightarrow{\operatorname{div}} H^{-1}(\Omega) \longrightarrow 0$$

is also exact. Then by Corollary 2.5, we get from the commutative diagram (2.26) that

$$(2.28) \quad (\mathbf{H}_0(\operatorname{curl}, \Omega))' = \mathbf{H}^{-1}(\operatorname{div}, \Omega).$$

The inclusion $\mathbf{H}^{-1}(\operatorname{div}, \Omega) \subset (\mathbf{H}_0(\operatorname{curl}, \Omega))'$ has been proved in the book [9, p. 338] for $n = 2$.

As we mentioned, $J_{H(\operatorname{curl})}$ is just an identity operator on $\nabla H_0^1(\Omega)$; thus the decomposition (2.27) is orthogonal in the inner product $(\cdot, \cdot)_{H(\operatorname{curl})'}$, but not in the L^2 inner product or in the H^{-1} inner product.

2.3. Helmholtz decomposition of $H^{-1}(\operatorname{curl})$ space. Following [23], we introduce the space

$$K_0^c := \{\phi \in \mathbf{H}_0(\operatorname{curl}, \Omega) : \operatorname{div} \phi = 0\} = \mathbf{H}_0(\operatorname{curl}, \Omega) / \operatorname{grad} H_0^1$$

equipped with norm $\|\cdot\|_{H(\operatorname{curl})}$. Noting that $\operatorname{curl} K_0^c = \operatorname{curl} \mathbf{H}_0(\operatorname{curl}, \Omega)$, we get the following exact sequence from the 3D de Rham complex (2.21):

$$0 \longrightarrow K_0^c \xrightarrow{\operatorname{curl}} \mathbf{H}_0(\operatorname{div}, \Omega) \xrightarrow{\operatorname{div}} L_0^2(\Omega) \longrightarrow 0,$$

which together with Remark 2.15 in [46] implies the exactness of the dual complex

$$0 \longrightarrow L_0^2(\Omega) \xrightarrow{\operatorname{grad}} (\mathbf{H}_0(\operatorname{div}, \Omega))' \xrightarrow{\operatorname{curl}} (K_0^c)' \longrightarrow 0.$$

We then construct an exact sequence with dual spaces to characterize the dual space $(\mathbf{H}_0(\operatorname{div}, \Omega))'$. Define

$$\mathbf{H}^{-1}(\operatorname{curl}, \Omega) = \{\phi \in \mathbf{H}^{-1}(\Omega; \mathbb{R}^3) : \operatorname{curl} \phi \in \mathbf{H}^{-1}(\Omega; \mathbb{R}^3)\}$$

with squared norm $\|\phi\|_{H^{-1}(\operatorname{curl})}^2 := \|\phi\|_{-1}^2 + \|\operatorname{curl} \phi\|_{-1}^2$.

LEMMA 2.7. *The complex*

$$0 \longrightarrow L_0^2(\Omega) \xrightarrow{\text{grad}} \mathbf{H}^{-1}(\text{curl}, \Omega) \xrightarrow{\text{curl}} (K_0^c)' \longrightarrow 0$$

is exact.

Proof. We have $\ker(\text{curl}) \cap \mathbf{H}^{-1}(\Omega; \mathbb{R}^3) = \text{grad } L_0^2(\Omega)$ by taking $s = -3$ in the exact sequence (2.24). Apparently $\ker(\text{curl}) \cap \mathbf{H}^{-1}(\text{curl}, \Omega) = \ker(\text{curl}) \cap \mathbf{H}^{-1}(\Omega; \mathbb{R}^3)$, which implies the exactness of the former complex.

By the Poincaré inequality on K_0^c [44, 34], $(\text{curl} \cdot, \text{curl} \cdot)$ defines an inner product on K_0^c and $(\text{curl curl})^{-1} : (K_0^c)' \rightarrow K_0^c$ is an isomorphism. Given an $\mathbf{f} \in (K_0^c)'$, find $\mathbf{u} \in K_0^c$ such that $\text{curl curl } \mathbf{u} = \mathbf{f}$ in $(K_0^c)'$ is the Maxwell's equation with divergence-free constraint. Since $\text{curl } K_0^c \subset \mathbf{H}^{-1}(\text{curl}, \Omega)$, we obtain

$$(K_0^c)' = \text{curl curl } K_0^c \subset \text{curl } \mathbf{H}^{-1}(\text{curl}, \Omega).$$

Due to (2.28),

$$\text{curl } \mathbf{H}^{-1}(\text{curl}, \Omega) \subset \mathbf{H}^{-1}(\text{div}, \Omega) = (\mathbf{H}_0(\text{curl}, \Omega))' \subset (K_0^c)'.$$

Therefore,

$$(2.29) \quad (K_0^c)' = \text{curl } \mathbf{H}^{-1}(\text{curl}, \Omega) = (\mathbf{H}_0(\text{curl}, \Omega))' = \mathbf{H}^{-1}(\text{div}, \Omega),$$

as required. □

Then we construct the commutative diagram

$$(2.30) \quad \begin{array}{ccccc} L_0^2(\Omega) & \xrightarrow{\text{grad}} & \mathbf{H}^{-1}(\text{curl}, \Omega) & \xrightarrow{\text{curl}} & (K_0^c)' \longrightarrow 0 \\ & & \uparrow I & & \uparrow \text{curl curl} \\ & & \mathbf{H}_0(\text{div}, \Omega) & \xleftarrow{\text{curl}} & K_0^c \end{array} \quad .$$

Using Theorem 2.2, the following stable Helmholtz decomposition holds:

$$(2.31) \quad \mathbf{H}^{-1}(\text{curl}, \Omega) = \nabla L_0^2(\Omega) \oplus \text{curl } K_0^c = \nabla L_0^2(\Omega) \oplus \text{curl } \mathbf{H}_0(\text{curl}, \Omega).$$

According to Corollary 2.5 and commutative diagram (2.30), it follows that

$$(\mathbf{H}_0(\text{div}, \Omega))' = \mathbf{H}^{-1}(\text{curl}, \Omega),$$

and thus the decomposition (2.31) is also orthogonal in the $(\cdot, \cdot)_{\mathbf{H}(\text{div})'}$ inner product. A decomposition of the dual space of $\mathbf{H}_0^m(\text{div}, \Omega) := \{\phi \in \mathbf{H}_0(\text{div}, \Omega) : \text{div } \phi \in H_0^m(\Omega)\}$ was presented in [2].

2.4. Helmholtz decomposition of symmetric tensors: HHJ complex.

We now consider differential complexes involving symmetric tensor functions.

LEMMA 2.8. *We have the following exact sequence:*

$$(2.32) \quad \mathbf{H}^1(\Omega; \mathbb{R}^2) \xrightarrow{\text{sym curl}} L^2(\Omega; \mathbb{S}) \xrightarrow{\text{div div}} H^{-2}(\Omega) \longrightarrow 0.$$

Proof. The identity $\ker(\operatorname{div} \mathbf{div}) = \operatorname{img}(\operatorname{sym} \operatorname{curl})$ can be found in [37]. Thanks to the exact sequence (2.20) with $s = -1, -2$,

$$\operatorname{div} \operatorname{div} \mathbf{L}^2(\Omega; \mathbb{M}) = \operatorname{div} \mathbf{H}^{-1}(\Omega; \mathbb{R}^2) = H^{-2}(\Omega).$$

Noting that $\mathbf{L}^2(\Omega; \mathbb{M}) = \mathbf{L}^2(\Omega; \mathbb{S}) + \mathbf{L}^2(\Omega; \mathbb{K})$ and $\operatorname{div} \operatorname{div} \mathbf{L}^2(\Omega; \mathbb{K}) = 0$, we achieve $\operatorname{div} \operatorname{div} \mathbf{L}^2(\Omega; \mathbb{S}) = \operatorname{div} \operatorname{div} \mathbf{L}^2(\Omega; \mathbb{M}) = H^{-2}(\Omega)$. \square

With the exact sequence (2.32) we construct the following commutative diagram:

$$\begin{array}{ccccc} \mathbf{H}^1(\Omega; \mathbb{R}^2) & \xrightarrow{\operatorname{sym} \operatorname{curl}} & \mathbf{L}^2(\Omega; \mathbb{S}) & \xrightarrow{\operatorname{div} \operatorname{div}} & H^{-2}(\Omega) & \longrightarrow & 0 \\ & & \uparrow I & & \uparrow \Delta^2 & & \\ & & \mathbf{L}^2(\Omega; \mathbb{S}) & \xleftarrow{\nabla^2} & H_0^2(\Omega) & & \end{array} .$$

By Corollary 2.4, we recover the L^2 -orthogonal Helmholtz decomposition obtained in [37, Lemma 3.1]:

$$\mathbf{L}^2(\Omega; \mathbb{S}) = \operatorname{sym} \operatorname{curl} \mathbf{H}^1(\Omega; \mathbb{R}^2) \oplus^\perp \nabla^2 H_0^2(\Omega).$$

We can follow the definition of $\mathbf{H}^{-1}(\operatorname{div} \operatorname{div}, \Omega)$ to introduce the following spaces:

$$\mathbf{H}^{-1}(\operatorname{div} \operatorname{div}, \Omega; \mathbb{S}) := \{ \boldsymbol{\tau} \in \mathbf{L}^2(\Omega; \mathbb{S}) : \operatorname{div} \operatorname{div} \boldsymbol{\tau} \in H^{-1}(\Omega) \}$$

with squared norm $\| \boldsymbol{\tau} \|_{\mathbf{H}^{-1}(\operatorname{div} \operatorname{div})}^2 := \| \boldsymbol{\tau} \|_0^2 + \| \operatorname{div} \operatorname{div} \boldsymbol{\tau} \|_{-1}^2$, and

$$\mathbf{H}(\operatorname{div} \operatorname{div}, \Omega; \mathbb{S}) := \{ \boldsymbol{\tau} \in \mathbf{L}^2(\Omega; \mathbb{S}) : \operatorname{div} \operatorname{div} \boldsymbol{\tau} \in L^2(\Omega) \}$$

with squared norm $\| \boldsymbol{\tau} \|_{\mathbf{H}(\operatorname{div} \operatorname{div})}^2 := \| \boldsymbol{\tau} \|_0^2 + \| \operatorname{div} \operatorname{div} \boldsymbol{\tau} \|_0^2$. We recall the Hellan–Herrmann–Johnson (HHJ) exact sequence (cf. [21, Lemma 2.2]) and give a simple proof here.

LEMMA 2.9. *We have the following exact sequence:*

$$(2.33) \quad \mathbf{H}^1(\Omega; \mathbb{R}^2) \xrightarrow{\operatorname{sym} \operatorname{curl}} \mathbf{H}^{-1}(\operatorname{div} \operatorname{div}, \Omega; \mathbb{S}) \xrightarrow{\operatorname{div} \operatorname{div}} H^{-1}(\Omega) \longrightarrow 0.$$

Proof. We only need to prove $\operatorname{div} \operatorname{div} \mathbf{H}^{-1}(\operatorname{div} \operatorname{div}, \Omega; \mathbb{S}) = H^{-1}(\Omega)$. By the definition of $\mathbf{H}^{-1}(\operatorname{div} \operatorname{div}, \Omega; \mathbb{S})$, apparently $\operatorname{div} \operatorname{div} \mathbf{H}^{-1}(\operatorname{div} \operatorname{div}, \Omega; \mathbb{S}) \subset H^{-1}(\Omega)$. On the other side, for each $v \in H^{-1}(\Omega) \subset H^{-2}(\Omega)$, by the exact sequence (2.32), there exists $\boldsymbol{\tau} \in \mathbf{L}^2(\Omega; \mathbb{S})$ such that $\operatorname{div} \operatorname{div} \boldsymbol{\tau} = v$. Note that $v \in H^{-1}(\Omega)$; thus $\boldsymbol{\tau} \in \mathbf{H}^{-1}(\operatorname{div} \operatorname{div}, \Omega; \mathbb{S})$, which indicates $H^{-1}(\Omega) \subset \operatorname{div} \operatorname{div} \mathbf{H}^{-1}(\operatorname{div} \operatorname{div}, \Omega; \mathbb{S})$. \square

Given a scalar function v , we can embed it into the symmetric tensor space as $\boldsymbol{\pi}(v) = v \mathbf{I}_{2 \times 2}$. Since $\Delta v = \operatorname{div} \operatorname{div} \boldsymbol{\pi}(v)$, we have the commutative diagram in two dimensions

$$(2.34) \quad \begin{array}{ccccc} \mathbf{H}^1(\Omega; \mathbb{R}^2) & \xrightarrow{\operatorname{sym} \operatorname{curl}} & \mathbf{H}^{-1}(\operatorname{div} \operatorname{div}, \Omega; \mathbb{S}) & \xrightarrow{\operatorname{div} \operatorname{div}} & H^{-1}(\Omega) & \longrightarrow & 0 \\ & & \uparrow I & & \uparrow \Delta & & \\ & & \mathbf{H}_0^1(\Omega; \mathbb{S}) & \xleftarrow{\boldsymbol{\pi}} & H_0^1(\Omega) & & \end{array} .$$

According to Theorem 2.2, we recover the recent Hemholtz decomposition in [40, Theorem 3.1]:

$$(2.35) \quad \mathbf{H}^{-1}(\operatorname{div} \mathbf{div}, \Omega; \mathbb{S}) = \operatorname{sym} \operatorname{curl} \mathbf{H}^1(\Omega; \mathbb{R}^2) \oplus \pi H_0^1(\Omega).$$

The index -1 can be further decreased to -2 ; see Lemma 2.10 in the next subsection. The generalized Hemholtz decomposition (2.35) is not orthogonal in $\mathbf{H}^{-1}(\Omega; \mathbb{S})$ or the $\mathbf{H}^{-1}(\operatorname{div} \mathbf{div}, \Omega; \mathbb{S})$ inner product. Indeed, $\pi \neq (\operatorname{div} \mathbf{div})'$; thus (2.35) is derived from Theorem 2.2 for Banach spaces rather than Corollary 2.4 for Hilbert spaces.

Smoothness of the symmetric tensor can be further increased to

$$\begin{array}{ccccc} \mathbf{H}^1(\Omega; \mathbb{R}^2) & \xrightarrow{\operatorname{sym} \operatorname{curl}} & \mathbf{H}(\operatorname{div} \mathbf{div}, \Omega; \mathbb{S}) & \xrightarrow{\operatorname{div} \mathbf{div}} & L^2(\Omega) & \longrightarrow & 0 \\ & & \uparrow I & & \uparrow I & & \\ & & \mathbf{H}_0^1(\Omega; \mathbb{S}) & \xleftarrow{\pi \Delta^{-1}} & L^2(\Omega) & & \end{array},$$

which leads to the generalized Helmholtz decomposition

$$\mathbf{H}(\operatorname{div} \mathbf{div}, \Omega; \mathbb{S}) = \operatorname{sym} \operatorname{curl} \mathbf{H}^1(\Omega; \mathbb{R}^2) \oplus \pi \Delta^{-1} L^2(\Omega).$$

2.5. Helmholtz decomposition of symmetric tensors: Linear elasticity.

Recall that the symmetric gradient $\varepsilon(\mathbf{u}) = (\nabla u + (\nabla u)^\top)/2$. Let

$$\mathbf{H}^{-2}(\operatorname{rot} \operatorname{rot}, \Omega; \mathbb{S}) := \{\boldsymbol{\tau} \in \mathbf{H}^{-1}(\Omega; \mathbb{S}) : \operatorname{rot} \operatorname{rot} \boldsymbol{\tau} \in H^{-2}(\Omega)\}$$

with squared norm $\|\boldsymbol{\tau}\|_{\mathbf{H}^{-2}(\operatorname{rot} \operatorname{rot})}^2 := \|\boldsymbol{\tau}\|_{-1}^2 + \|\operatorname{rot} \operatorname{rot} \boldsymbol{\tau}\|_{-2}^2$.

LEMMA 2.10. *The complex*

$$(2.36) \quad L^2(\Omega; \mathbb{R}^2) \xrightarrow{\varepsilon} \mathbf{H}^{-2}(\operatorname{rot} \operatorname{rot}, \Omega; \mathbb{S}) \xrightarrow{\operatorname{rot} \operatorname{rot}} H^{-2}(\Omega) \longrightarrow 0$$

is exact.

Proof. It is trivial that (2.36) is a complex, i.e., $\operatorname{rot} \operatorname{rot} \circ \varepsilon = 0$. Next we show the exactness. For any $\boldsymbol{\tau} \in \ker(\operatorname{rot} \operatorname{rot})$, by the exact sequence (2.25) with $s = -3$, there exists $v \in H^{-1}(\Omega)$ satisfying $\operatorname{rot} \boldsymbol{\tau} = \nabla v$. Since $\nabla v = \operatorname{rot} \begin{pmatrix} 0 & v \\ -v & 0 \end{pmatrix}$, we have

$$\operatorname{rot} \left(\boldsymbol{\tau} - \begin{pmatrix} 0 & v \\ -v & 0 \end{pmatrix} \right) = \mathbf{0}.$$

Thus $\boldsymbol{\tau} = \begin{pmatrix} 0 & v \\ -v & 0 \end{pmatrix} + \nabla \phi$ with $\phi \in L^2(\Omega; \mathbb{R}^2)$, which together with the fact that $\boldsymbol{\tau}$ is symmetric means $\boldsymbol{\tau} = \varepsilon(\phi)$. Hence $\ker(\operatorname{rot} \operatorname{rot}) \subset \operatorname{img}(\varepsilon)$.

By the rotation of the exact sequence (2.32), $\operatorname{rot} \operatorname{rot} L^2(\Omega; \mathbb{S}) = H^{-2}(\Omega)$. Thus $H^{-2}(\Omega) \subset \operatorname{rot} \operatorname{rot} \mathbf{H}^{-2}(\operatorname{rot} \operatorname{rot}, \Omega; \mathbb{S})$, which implies $\operatorname{img}(\operatorname{rot} \operatorname{rot}) = H^{-2}(\Omega)$. \square

With the complex (2.36) and the fact that $\Delta^2 = \operatorname{rot} \operatorname{rot} \mathbf{I} \operatorname{curl} \operatorname{curl}$, we have the commutative diagram

$$(2.37) \quad \begin{array}{ccccc} L^2(\Omega; \mathbb{R}^2) & \xrightarrow{\varepsilon} & \mathbf{H}^{-2}(\operatorname{rot} \operatorname{rot}, \Omega; \mathbb{S}) & \xrightarrow{\operatorname{rot} \operatorname{rot}} & H^{-2}(\Omega) & \longrightarrow & 0 \\ & & \uparrow I & & \uparrow \Delta^2 & & \\ & & \mathbf{H}_0(\operatorname{div}, \Omega; \mathbb{S}) & \xleftarrow{\operatorname{curl} \operatorname{curl}} & H_0^2(\Omega) & & \end{array},$$

which leads to a Helmholtz decomposition:

$$(2.38) \quad \mathbf{H}^{-2}(\text{rot rot}, \Omega; \mathbb{S}) = \varepsilon \mathbf{L}^2(\Omega; \mathbb{R}^2) \oplus \mathbf{curl curl} H_0^2(\Omega).$$

Applying Corollary 2.5 to the commutative diagram (2.37), we have

$$(\mathbf{H}_0(\text{div}, \Omega; \mathbb{S}))' = \mathbf{H}^{-2}(\text{rot rot}, \Omega; \mathbb{S}).$$

Therefore, the decomposition (2.38) is also orthogonal in the $\mathbf{H}_0(\text{div}, \Omega; \mathbb{S})'$ inner product.

2.6. Helmholtz decomposition of $\mathbf{H}(\text{curl curl}, (K_0^c)')$ space. Let

$$\mathbf{L}_0^2(\Omega; \mathbb{M}) := \{\boldsymbol{\tau} \in \mathbf{L}^2(\Omega; \mathbb{M}) : (\text{tr } \boldsymbol{\tau}, 1) = 0\}.$$

Introduce two Hilbert spaces

$$\mathbf{H}^0(\mathbf{div}, \text{dev sym}) := \{\boldsymbol{\tau} \in \mathbf{L}_0^2(\Omega; \mathbb{M}) : \text{div } \boldsymbol{\tau} = \mathbf{0}, \text{dev sym } \boldsymbol{\tau} = \mathbf{0}\}$$

with norm $\|\cdot\|_0$, and

$$\mathbf{H}(\text{curl curl}, (K_0^c)') := \{\boldsymbol{\phi} \in \mathbf{L}^2(\Omega; \mathbb{R}^3) : \text{curl curl } \boldsymbol{\phi} \in (K_0^c)'\}$$

with squared norm $\|\boldsymbol{\tau}\|_{\mathbf{H}(\text{curl curl}, (K_0^c)')}^2 := \|\boldsymbol{\tau}\|_0^2 + \|\text{curl curl } \boldsymbol{\tau}\|_{(K_0^c)'}^2$.

LEMMA 2.11. *The complex*

$$\mathbf{H}^0(\mathbf{div}, \text{dev sym}) \xrightarrow{\text{spn}^{-1} \text{skw}} \mathbf{H}(\text{curl curl}, (K_0^c)') \xrightarrow{\text{curl curl}} (K_0^c)' \longrightarrow 0$$

is exact.

Proof. By definition $\text{img}(\text{curl curl}) = (K_0^c)'$. Next we show that $\ker(\text{curl curl}) = \text{img}(\text{spn}^{-1} \text{skw})$.

Taking any $\boldsymbol{\tau} \in \mathbf{H}^0(\mathbf{div}, \text{dev sym})$, set $\boldsymbol{\psi} = \text{spn}^{-1} \text{skw } \boldsymbol{\tau}$. Then we have $\text{spn } \boldsymbol{\psi} = \text{skw } \boldsymbol{\tau}$. Noting that $\text{dev sym } \boldsymbol{\tau} = \mathbf{0}$, i.e., $\text{sym } \boldsymbol{\tau} = \frac{1}{3}(\text{tr } \boldsymbol{\tau})\mathbf{I}$, it holds that

$$\boldsymbol{\tau} = \text{sym } \boldsymbol{\tau} + \text{skw } \boldsymbol{\tau} = \frac{1}{3}(\text{tr } \boldsymbol{\tau})\mathbf{I} + \text{spn } \boldsymbol{\psi}.$$

Hence by $\text{div } \boldsymbol{\tau} = \mathbf{0}$ we obtain $\frac{1}{3}\nabla(\text{tr } \boldsymbol{\tau}) = -\mathbf{div} \text{spn } \boldsymbol{\psi} = \text{curl } \boldsymbol{\psi}$, which implies $\text{curl curl } \boldsymbol{\psi} = \mathbf{0}$. Thus we have $\text{img}(\text{spn}^{-1} \text{skw}) \subset \ker(\text{curl curl})$.

On the other hand, take any $\boldsymbol{\psi} \in \ker(\text{curl curl})$. Then there exists $w \in L_0^2(\Omega)$ such that $\text{curl } \boldsymbol{\psi} = \nabla w$. Let

$$(2.39) \quad \boldsymbol{\tau} = \text{spn } \boldsymbol{\psi} + w\mathbf{I}.$$

Obviously $\mathbf{div} \boldsymbol{\tau} = 0$, $w\mathbf{I} = \text{sym } \boldsymbol{\tau}$, and $w = \frac{1}{3} \text{tr}(\text{sym } \boldsymbol{\tau})$. Thus $\text{dev sym } \boldsymbol{\tau} = \mathbf{0}$; i.e., $\boldsymbol{\tau} \in \mathbf{H}^0(\mathbf{div}, \text{dev sym})$. By applying skw to (2.39), we have $\text{spn } \boldsymbol{\psi} = \text{skw } \boldsymbol{\tau}$, which indicates that $\boldsymbol{\psi} = \text{spn}^{-1} \text{skw } \boldsymbol{\tau} \in \text{img}(\text{spn}^{-1} \text{skw})$. □

According to the proof of Lemma 2.11, we have

$$(2.40) \quad \mathbf{H}^0(\mathbf{div}, \text{dev sym}) = \{\boldsymbol{\tau} = v\mathbf{I} + \text{spn } \boldsymbol{\phi} : v \in L_0^2(\Omega), \boldsymbol{\phi} \in \mathbf{L}^2(\Omega; \mathbb{R}^3), \text{div } \boldsymbol{\tau} = 0\}.$$

Therefore, we have the commutative diagram

$$(2.41) \quad \begin{array}{ccccccc} \mathbf{H}^0(\mathbf{div}, \text{dev sym}) & \xrightarrow{\text{spn}^{-1} \text{skw}} & \mathbf{H}(\text{curl curl}, (K_0^c)') & \xrightarrow{\text{curl curl}} & (K_0^c)' & \longrightarrow & 0 \\ & & \uparrow I & & \uparrow \text{curl curl} & & \\ & & K_0^c & \xleftarrow{I} & K_0^c & & \end{array} .$$

Applying Theorem 2.2, we get the stable Helmholtz decomposition

$$(2.42) \quad \mathbf{H}(\text{curl curl}, (K_0^c)') = \text{spn}^{-1} \text{skw } \mathbf{H}^0(\mathbf{div}, \text{dev sym}) \oplus K_0^c.$$

Again, this Helmholtz decomposition is not orthogonal in the $\mathbf{H}(\text{curl curl}, (K_0^c)')$ or $(K_0^c)'$ inner product.

2.7. Helmholtz decomposition of $H^{-2}(\text{div}^3, \Omega)$ space. Denote

$$\mathbf{H}^{-2}(\text{div}^3, \Omega) := \{ \boldsymbol{\tau} \in \mathbf{L}^2(\Omega; \mathbb{S}(3)) : \text{div div div } \boldsymbol{\tau} \in H^{-2}(\Omega) \}$$

with squared norm $\| \boldsymbol{\tau} \|_{\mathbf{H}^{-2}(\text{div}^3)}^2 := \| \boldsymbol{\tau} \|_0^2 + \| \text{div div div } \boldsymbol{\tau} \|_{-2}^2$. Define $\Xi : H_0^1(\Omega; \mathbb{R}^2) \rightarrow \mathbf{H}^{-2}(\text{div}^3, \Omega)$ as follows: for any $\boldsymbol{\psi} = (\psi_1, \psi_2)^\top \in H_0^1(\Omega; \mathbb{R}^2)$, $\Xi \boldsymbol{\psi} := (\boldsymbol{\tau}_{ijk})_{2 \times 2 \times 2}$ with

$$\boldsymbol{\tau}_{111} = \psi_1, \boldsymbol{\tau}_{222} = \psi_2, \boldsymbol{\tau}_{112} = \boldsymbol{\tau}_{121} = \boldsymbol{\tau}_{211} = \frac{1}{3} \psi_2, \boldsymbol{\tau}_{122} = \boldsymbol{\tau}_{212} = \boldsymbol{\tau}_{221} = \frac{1}{3} \psi_1.$$

LEMMA 2.12. *The complex*

$$\mathbf{H}^1(\Omega; \mathbb{S}) \xrightarrow{\text{sym curl}} \mathbf{H}^{-2}(\text{div}^3, \Omega) \xrightarrow{\text{div div div}} H^{-2}(\Omega) \longrightarrow 0$$

is exact.

Proof. It is apparent that $\Delta^2 = \text{div div div } \Xi \nabla$; thus $\text{img}(\text{div div div}) = H^{-2}(\Omega)$. We refer the reader to [48, Theorem 4.1 and Theorem 4.4] for $\ker(\text{div div div}) = \text{img}(\text{sym curl})$. \square

Through constructing the commutative diagram

$$(2.43) \quad \begin{array}{ccccccc} \mathbf{H}^1(\Omega; \mathbb{S}) & \xrightarrow{\text{sym curl}} & \mathbf{H}^{-2}(\text{div}^3, \Omega) & \xrightarrow{\text{div div div}} & H^{-2}(\Omega) & \longrightarrow & 0 \\ & & \Xi \uparrow & & \uparrow \Delta^2 & & \\ & & H_0^1(\Omega; \mathbb{R}^2) & \xleftarrow{\nabla} & H_0^2(\Omega) & & \end{array} ,$$

we get the stable Helmholtz decomposition from Theorem 2.2:

$$(2.44) \quad \mathbf{H}^{-2}(\text{div}^3, \Omega) = \text{sym curl } \mathbf{H}^1(\Omega; \mathbb{S}) \oplus \Xi \nabla H_0^2(\Omega).$$

More differential complexes and Helmholtz decompositions can be obtained, and some of them will be discussed along with the mixed formulations of elliptic systems.

3. Abstract mixed formulation and its decomposition. In this section we present an abstract mixed formulation and use a Helmholtz decomposition to decouple the saddle point system into several elliptic problems.

3.1. Framework. Assume we have the exact sequence

$$(3.1) \quad P \xrightarrow{d^-} \Sigma \xrightarrow{d} V' \longrightarrow 0$$

and the commutative diagram

$$(3.2) \quad \begin{array}{ccccccc} & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ X & \xrightarrow{J_X} & X' & & & & \\ & & \cup & & & & \\ P & \xrightarrow{d^-} & \Sigma & \xrightarrow{d} & V' & \longrightarrow & 0 \\ & & \uparrow \Pi_\Sigma & & \uparrow J_V & & \\ & & \tilde{\Sigma} & \xleftarrow{\Pi_V} & V & & \end{array} ,$$

where the isomorphisms J_X and J_V are given by (2.7), i.e., the inverse of the Riesz representation operator, and the rest of the linear operators are all continuous but not necessarily isomorphic.

By Theorem 2.2, we have a stable Helmholtz decomposition:

$$(3.3) \quad \Sigma = d^- P \oplus \Pi_\Sigma \Pi_V V.$$

We emphasize that we do not need to know either the short exact sequence at the bottom or the space $\tilde{\Sigma}$ in a very precise form (i.e., $\tilde{\Sigma}$ can be reasonably enlarged to include the image space $\Pi_V V$).

3.1.1. Two-term formulation. An abstract mixed formulation based on the commutative diagram (3.2) is as follows: given $g \in \Sigma'$ and $f \in V'$, find $(\sigma, u) \in \Sigma \times V$ such that

$$(3.4) \quad (\sigma, \tau)_{X'} + \langle d\tau, u \rangle = \langle g, \tau \rangle \quad \forall \tau \in \Sigma,$$

$$(3.5) \quad \langle d\sigma, v \rangle = \langle f, v \rangle \quad \forall v \in V.$$

The bilinear form $(\cdot, \cdot)_{X'}$ does not need to be an inner product unless we intend to involve X and J_X in the mixed formulation. We only require that $(\cdot, \cdot)_{X'}$ be positive semidefinite and symmetric.

To show the well-posedness of the mixed formulation (3.4)–(3.5), we assume the following norm equivalence:

$$(3.6) \quad \|\tau\|_\Sigma^2 \approx \|\tau\|_{X'}^2 + \|d\tau\|_{V'}^2, \quad \forall \tau \in \Sigma.$$

This norm equivalence is usually trivial, and it holds apparently for all the examples in this paper. Indeed, the space Σ is usually constructed so that (3.6) holds.

THEOREM 3.1. Assume the exact sequence (3.1), the commutative diagram (3.2), and the norm equivalence (3.6) hold; then the mixed formulation (3.4)–(3.5) is uniquely solvable. Moreover, we have the stability result

$$\|\sigma\|_\Sigma + \|u\|_V \lesssim \|g\|_{\Sigma'} + \|f\|_{V'}.$$

Proof. It is trivial that the bilinear forms in the mixed formulation (3.4)–(3.5) are continuous due to (3.6). Using (3.6) again, it is also obvious that

$$\|\tau\|_\Sigma \lesssim \|\tau\|_{X'} + \|d\tau\|_{V'} = \|\tau\|_{X'} \quad \forall \tau \in \ker d.$$

By the Babuška–Brezzi theory (cf. [6, 13, 8]), it suffices to prove the inf-sup condition

$$(3.7) \quad \|v\|_V \lesssim \sup_{\tau \in \Sigma} \frac{\langle d\tau, v \rangle}{\|\tau\|_\Sigma} \quad \forall v \in V.$$

For each $v \in V$, let $\tau = \Pi_\Sigma \Pi_V v$. It is apparent that

$$\|\tau\|_\Sigma = \|\Pi_\Sigma \Pi_V v\|_\Sigma \lesssim \|v\|_V.$$

Then, making use of the commutative diagram (3.2) and (2.7), it follows that

$$\langle d\tau, v \rangle = \langle d\Pi_\Sigma \Pi_V v, v \rangle = \langle J_V v, v \rangle = \|v\|_V^2.$$

Hence we have

$$\|v\|_V \|\tau\|_\Sigma \lesssim \|v\|_V^2 = \langle d\tau, v \rangle,$$

which means the inf-sup condition (3.7). □

3.1.2. Unfolded three-term formulation. We derive an equivalent three-term formulation of the mixed formulation (3.4)–(3.5) when X' is a Sobolev space of negative order. In this subsection, we assume the bilinear form $(\cdot, \cdot)_{X'}$ is the corresponding inner product of X' .

Let $\phi = J_X^{-1} \sigma \in X$. By (2.8), we can rewrite (3.4) as

$$\langle \tau, \phi \rangle + \langle d\tau, u \rangle = \langle g, \tau \rangle \quad \forall \tau \in \Sigma.$$

Noting that $\sigma = J_X \phi$, it follows from (2.7) that

$$\langle \sigma, \psi \rangle = \langle J_X \phi, \psi \rangle = (\phi, \psi)_X \quad \forall \psi \in X.$$

Therefore, the mixed formulation (3.4)–(3.5) is equivalent to an unfolded three-term formulation: find $(\phi, u, \sigma) \in X \times V \times \Sigma$ such that

$$(3.8) \quad (\phi, \psi)_X - \langle \sigma, d'v + \psi \rangle = -\langle f, v \rangle + \langle g_X, \psi \rangle \quad \forall (\psi, v) \in X \times V,$$

$$(3.9) \quad \langle d'u + \phi, \tau \rangle = \langle g, \tau \rangle \quad \forall \tau \in \Sigma,$$

with $g_X = 0$. It is interesting to note that the variable σ can be formally interpreted as the Lagrange multiplier to impose the constraint $I'\phi = -d'u$ in Σ' if $g = 0$, where $I : \Sigma \rightarrow X'$ is the natural embedding operator. The operator equation of the mixed formulation (3.8)–(3.9) is

$$(3.10) \quad \begin{pmatrix} J_X & 0 & -I \\ 0 & 0 & -d \\ I' & d' & 0 \end{pmatrix} \begin{pmatrix} \phi \\ u \\ \sigma \end{pmatrix} = \begin{pmatrix} g_X \\ -f \\ g \end{pmatrix}.$$

According to Theorem 3.1, we immediately obtain the well-posedness of the mixed formulation (3.8)–(3.9) for $g_X = 0$. The well-posedness of the mixed formulation (3.8)–(3.9) for general $g_X \in X'$ is given as follows.

THEOREM 3.2. *Assume the exact sequence (3.1), the commutative diagram (3.2), and the norm equivalence (3.6) hold; then the unfolded mixed formulation (3.8)–(3.9) is uniquely solvable. Moreover, we have the stability result*

$$\|\phi\|_X + \|\sigma\|_\Sigma + \|u\|_V \lesssim \|g_X\|_{X'} + \|g\|_{\Sigma'} + \|f\|_{V'}.$$

Proof. We split (3.10) as a 2×2 block saddle point system by treating $X \times V$ as one space and denote $B = [I' \ d'] : X \times V \rightarrow \Sigma'$ as $B((\phi, u)) = I'\phi + d'u$.

By (3.6), all the bilinear forms in the mixed formulation (3.8)–(3.9) are obviously continuous. Let $(\psi, v) \in \ker(B) \cap (X \times V)$, i.e., $I'\psi = -d'v$. Due to (2.7) and the commutative diagram (3.2), we get

$$\begin{aligned} \|v\|_V^2 &= \langle J_V v, v \rangle = \langle d\Pi_\Sigma \Pi_V v, v \rangle = \langle \Pi_\Sigma \Pi_V v, d'v \rangle \\ &\leq \| \Pi_\Sigma \Pi_V v \|_\Sigma \| d'v \|_{\Sigma'} \lesssim \|v\|_V \|d'v\|_{\Sigma'}. \end{aligned}$$

Noting that $I' : X \rightarrow \Sigma'$ is continuous, we have

$$\|v\|_V \lesssim \|d'v\|_{\Sigma'} = \|I'\psi\|_{\Sigma'} \lesssim \|\psi\|_X,$$

which implies the coercivity on the kernel of B .

On the other hand, for any $\tau \in \Sigma$ it follows from (3.6) that

$$\|\tau\|_\Sigma \lesssim \|\tau\|_{X'} + \|d\tau\|_{V'} = \sup_{\psi \in X} \frac{\langle \tau, I'\psi \rangle}{\|\psi\|_X} + \sup_{v \in V} \frac{\langle d\tau, v \rangle}{\|v\|_V} \lesssim \sup_{\psi \in X, v \in V} \frac{\langle \tau, I'\psi + d'v \rangle}{\|\psi\|_X + \|v\|_V},$$

which is just the inf-sup condition of B . Therefore, the required result is guaranteed by the Babuška–Brezzi theory. \square

3.1.3. Decoupled formulation. We decompose the mixed formulation (3.4)–(3.5) using the Helmholtz decomposition (3.3). We apply the Helmholtz decomposition (3.3) to both the trial and test functions

$$\sigma = d^-p + \Pi_\Sigma \Pi_V w, \quad \tau = d^-q + \Pi_\Sigma \Pi_V \chi,$$

where $p, q \in P/\ker d^-$, and $w, \chi \in V$. Then, substituting them into the mixed formulation (3.4)–(3.5), we have

$$(3.11) \quad (d^-p + \Pi_\Sigma \Pi_V w, \Pi_\Sigma \Pi_V \chi)_{X'} + \langle d\Pi_\Sigma \Pi_V \chi, u \rangle = \langle g, \Pi_\Sigma \Pi_V \chi \rangle,$$

$$(3.12) \quad (d^-p + \Pi_\Sigma \Pi_V w, d^-q)_{X'} = \langle g, d^-q \rangle,$$

$$(3.13) \quad \langle d\Pi_\Sigma \Pi_V w, v \rangle = \langle f, v \rangle$$

for any $\chi \in V$, $q \in P/\ker d^-$, and $v \in V$. We obtain from the commutative diagram (3.2) and (2.7) again

$$\langle d\Pi_\Sigma \Pi_V w, v \rangle = \langle J_V w, v \rangle = (w, v)_V, \quad \langle d\Pi_\Sigma \Pi_V \chi, u \rangle = (\chi, u)_V.$$

Therefore, the mixed formulation (3.11)–(3.13) is equivalent to (in backwards order) the following: find $w \in V$, $p \in P/\ker d^-$, and $u \in V$ such that

$$(3.14) \quad (w, v)_V = \langle f, v \rangle \quad \forall v \in V,$$

$$(3.15) \quad (d^-p, d^-q)_{X'} = \langle g, d^-q \rangle - (\Pi_\Sigma \Pi_V w, d^-q)_{X'} \quad \forall q \in P/\ker d^-,$$

$$(3.16) \quad (u, \chi)_V = \langle g, \Pi_\Sigma \Pi_V \chi \rangle - (\sigma, \Pi_\Sigma \Pi_V \chi)_{X'} \quad \forall \chi \in V,$$

where $\sigma = d^-p + \Pi_\Sigma \Pi_V w$.

Remark 3.3. When the decomposition (3.3) is orthogonal with respect to $(\cdot, \cdot)_{X'}$ and $g = 0$, the second equation (3.15) will disappear.

Applying the Helmholtz decomposition (3.3) to the unfolded formulation, the decoupled mixed formulation (3.8)–(3.9) is equivalent to finding $w, u \in V$, $\phi \in X$, and $p \in P/\ker d^-$ such that

$$\begin{aligned}
 (3.17) \quad & (w, v)_V = \langle f, v \rangle \quad \forall v \in V, \\
 (3.18) \quad & (\phi, \psi)_X - \langle d^- p, \psi \rangle = \langle \Pi_\Sigma \Pi_V w, \psi \rangle \quad \forall \psi \in X, \\
 (3.19) \quad & \langle d^- q, \phi \rangle = \langle g, d^- q \rangle \quad \forall q \in P/\ker d^-, \\
 (3.20) \quad & (u, \chi)_V = \langle g - \phi, \Pi_\Sigma \Pi_V \chi \rangle \quad \forall \chi \in V.
 \end{aligned}$$

The middle system (3.18)–(3.19) of (ϕ, p) is now a Stokes-type system.

We summarize the former derivation as follows.

THEOREM 3.4. *Assume the exact sequence (3.1), the commutative diagram (3.2), and the norm equivalence (3.6) hold; then the mixed formulation (3.4)–(3.5) can be decoupled as three elliptic equations (3.14)–(3.16) or four equations (3.17)–(3.20).*

Remark 3.5. Let $(\cdot, \cdot)_\Sigma$ be the inner product of Hilbert space Σ . When the space of harmonic forms

$$\mathfrak{H} := \{ \tau \in \Sigma : d\tau = 0, (\tau, d^- q)_\Sigma = 0 \quad \forall q \in P \}$$

is nontrivial, by Remark 2.3 we have the Helmholtz decomposition

$$\Sigma = d^- P \oplus \Pi_\Sigma \Pi_V V \oplus \mathfrak{H}.$$

Then the mixed formulation (3.4)–(3.5) can be decoupled as follows: find $w, u \in V$, $p \in P/\ker d^-$, and $r \in \mathfrak{H}$ such that

$$\begin{aligned}
 & (w, v)_V = \langle f, v \rangle \quad \forall v \in V, \\
 & (d^- p + r, d^- q + s)_{X'} = \langle g, d^- q + s \rangle - (\Pi_\Sigma \Pi_V w, d^- q + s)_{X'} \quad \forall q \in P/\ker d^-, s \in \mathfrak{H}, \\
 & (u, \chi)_V = \langle g, \Pi_\Sigma \Pi_V \chi \rangle - (\sigma, \Pi_\Sigma \Pi_V \chi)_{X'} \quad \forall \chi \in V,
 \end{aligned}$$

where $\sigma = d^- p + \Pi_\Sigma \Pi_V w + r$. A similar modification can be applied to the decoupling of the three-term formulation. In the decoupled formulation, however, the space of harmonic forms should be identified a priori, which may not be easy for complicated geometric domains.

In the rest of this section, we shall apply our abstract framework to several concrete examples. It is worth mentioning again that the norm equivalence (3.6) is trivial for all examples except the primal formulation of the fourth order curl equation. For this exceptional case, the norm equivalence (3.6) is the result of (2.29). Hence we will focus on the derivation of the commutative diagrams.

3.2. The primal formulation of the biharmonic equation. Consider the biharmonic equation with homogeneous Dirichlet boundary condition

$$(3.21) \quad \begin{cases} \Delta^2 u = f & \text{in } \Omega, \\ u = \partial_\nu u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $f \in H^{-1}(\Omega)$ and $\Omega \subset \mathbb{R}^n$ with $n = 2, 3$. The primal formulation of (3.21) is to find $u \in H_0^2(\Omega)$ such that

$$(3.22) \quad (\nabla^2 u, \nabla^2 v) = \langle f, v \rangle \quad \forall v \in H_0^2(\Omega).$$

We first discuss the decoupling of (3.22) in two dimensions. We recall the known decoupling from [36, 38, 27], which fits into the framework of this paper as follows. By the rotated version of the commutative diagram (2.26), we build up the commutative diagram

$$\begin{array}{ccccccc}
 & & \mathbf{H}_0^1(\Omega; \mathbb{R}^2) & \xrightarrow{\Delta} & \mathbf{H}^{-1}(\Omega; \mathbb{R}^2) & & \\
 & & & & \cup & & \\
 (3.23) & L_0^2(\Omega) & \xrightarrow{\text{grad}} & \mathbf{H}^{-1}(\text{rot}, \Omega) & \xrightarrow{\text{rot}} & H^{-1}(\Omega) & \longrightarrow 0 \quad , \\
 & & & \uparrow \Gamma & & \uparrow \Delta & \\
 & & & \mathbf{H}_0(\text{div}, \Omega) & \xleftarrow{\text{curl}} & H_0^1(\Omega) &
 \end{array}$$

where we recall that

$$\mathbf{H}^{-1}(\text{rot}, \Omega) := \{ \phi \in \mathbf{H}^{-1}(\Omega; \mathbb{R}^2) : \text{rot } \phi \in H^{-1}(\Omega) \}$$

with norm $\|\phi\|_{\mathbf{H}^{-1}(\text{rot})}^2 := \|\phi\|_{-1}^2 + \|\text{rot } \phi\|_{-1}^2$. According to Theorem 2.2, we have the Helmholtz decomposition

$$(3.24) \quad \mathbf{H}^{-1}(\text{rot}, \Omega) = \nabla L_0^2(\Omega) \oplus \text{curl } H_0^1(\Omega).$$

The corresponding mixed formulation is to find $(\gamma, u) \in \mathbf{H}^{-1}(\text{rot}, \Omega) \times H_0^1(\Omega)$ such that

$$(3.25) \quad (\gamma, \beta)_{-1} - \langle \text{rot } \beta, u \rangle = 0 \quad \forall \beta \in \mathbf{H}^{-1}(\text{rot}, \Omega),$$

$$(3.26) \quad \langle \text{rot } \gamma, v \rangle = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega),$$

where $f \in H^{-1}(\Omega)$ and $(\gamma, \beta)_{-1} := -\langle \Delta^{-1} \gamma, \beta \rangle = -\langle \gamma, \Delta^{-1} \beta \rangle$.

By introducing variable $\phi = -\Delta^{-1} \gamma \in \mathbf{H}_0^1(\Omega; \mathbb{R}^2)$, the unfolded formulation is to find $(\gamma, u, \phi) \in \mathbf{H}^{-1}(\text{rot}, \Omega) \times H_0^1(\Omega) \times \mathbf{H}_0^1(\Omega; \mathbb{R}^2)$ such that

$$(3.27) \quad (\nabla \phi, \nabla \psi) + \langle \gamma, \text{curl } v - \psi \rangle = \langle f, v \rangle \quad \forall (v, \psi) \in H_0^1(\Omega) \times \mathbf{H}_0^1(\Omega; \mathbb{R}^2),$$

$$(3.28) \quad \langle \beta, \text{curl } u - \phi \rangle = 0 \quad \forall \beta \in \mathbf{H}^{-1}(\text{rot}, \Omega),$$

which is just the rotation form of problem (2.4) in [15].

Equation (3.28) implies $\phi = \text{curl } u$, which is plugged into (3.27) by taking $\psi = \text{curl } v$ with $v \in H_0^2(\Omega)$, which gives the primal formulation of the biharmonic equation (3.22) in two dimensions.

According to Theorem 3.4, the decoupled and unfolded formulation is to find $w \in H_0^1(\Omega)$, $\phi \in \mathbf{H}_0^1(\Omega; \mathbb{R}^2)$, $p \in L_0^2(\Omega)$, and $u \in H_0^1(\Omega)$ such that

$$(3.29) \quad (\text{curl } w, \text{curl } v) = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega),$$

$$(3.30) \quad (\nabla \phi, \nabla \psi) + (\text{div } \psi, p) = (\text{curl } w, \psi) \quad \forall \psi \in \mathbf{H}_0^1(\Omega; \mathbb{R}^2),$$

$$(3.31) \quad (\text{div } \phi, q) = 0 \quad \forall q \in L_0^2(\Omega),$$

$$(3.32) \quad (\text{curl } u, \text{curl } \chi) = (\phi, \text{curl } \chi) \quad \forall \chi \in H_0^1(\Omega).$$

Therefore, we recover the decoupling that the primal formulation of the biharmonic equation (3.22) in two dimensions is equivalent to two Poisson equations and one Stokes equation [36, 38, 27].

Such decoupling of the biharmonic equation in two dimensions can be generalized in various ways. First, we consider the fourth order elliptic singular perturbation problem with homogeneous Dirichlet boundary condition

$$(3.33) \quad \varepsilon^2 \Delta^2 u - \Delta u = f.$$

If we equip space $\mathbf{H}_0^1(\Omega; \mathbb{R}^2)$ with norm $(\|\cdot\|_0^2 + \varepsilon^2 |\cdot|_1^2)^{1/2}$ where $\varepsilon \geq 0$, the fourth order elliptic singular perturbation problem (3.33) will be decoupled to find $w \in H_0^1(\Omega)$, $\phi \in \mathbf{H}_0^1(\Omega; \mathbb{R}^2)$, $p \in L_0^2(\Omega)$, and $u \in H_0^1(\Omega)$ such that

$$\begin{aligned} (\operatorname{curl} w, \operatorname{curl} v) &= \langle f, v \rangle & \forall v \in H_0^1(\Omega), \\ (\phi, \psi) + \varepsilon^2 (\nabla \phi, \nabla \psi) + (\operatorname{div} \psi, p) &= (\operatorname{curl} w, \psi) & \forall \psi \in \mathbf{H}_0^1(\Omega; \mathbb{R}^2), \\ (\operatorname{div} \phi, q) &= 0 & \forall q \in L_0^2(\Omega), \\ (\operatorname{curl} u, \operatorname{curl} \chi) &= (\phi, \operatorname{curl} \chi) & \forall \chi \in H_0^1(\Omega). \end{aligned}$$

Note that we derived this decoupling in 2016 independently of [27]. The second and third equations form the Brinkman problem. The robust well-posedness of the Brinkman problem with respect to the parameter ε can be found in [43, 45, 41, 42, 50]. In [43, 45, 41], the norms of the spaces $\mathbf{L}^2(\Omega; \mathbb{R}^2) \cap \varepsilon \mathbf{H}_0^1(\Omega; \mathbb{R}^2)$ and $H^1(\Omega) \cap L_0^2(\Omega) + \varepsilon^{-1} L_0^2(\Omega)$ were adopted for ϕ and p , respectively, which induced an efficient and robust preconditioner.

We then discuss the decoupling of (3.22) in three dimensions. Based on the commutative diagram (2.26), the primal formulation (3.22) of the biharmonic equation in three dimensions is equivalent to finding $w \in H_0^1(\Omega)$, $\phi \in \mathbf{H}_0^1(\Omega; \mathbb{R}^3)$, $p \in \mathbf{L}^2(\Omega; \mathbb{R}^3)/\nabla H^1(\Omega)$, and $u \in H_0^1(\Omega)$ such that (cf. [27])

$$\begin{aligned} (\nabla w, \nabla v) &= \langle f, v \rangle & \forall v \in H_0^1(\Omega), \\ (\nabla \phi, \nabla \psi) + (\operatorname{curl} \psi, p) &= (\nabla w, \psi) & \forall \psi \in \mathbf{H}_0^1(\Omega; \mathbb{R}^3), \\ (\operatorname{curl} \phi, q) &= 0 & \forall q \in \mathbf{L}^2(\Omega; \mathbb{R}^3)/\nabla H^1(\Omega), \\ (\nabla u, \nabla \chi) &= (\phi, \nabla \chi) & \forall \chi \in H_0^1(\Omega). \end{aligned}$$

3.3. HHJ mixed formulation. The Hellan–Herrmann–Johnson (HHJ) mixed formulation [31, 32, 39] of the biharmonic equation (3.21) in two dimensions is to find $(\sigma, u) \in \mathbf{H}^{-1}(\operatorname{div} \operatorname{div}, \Omega; \mathbb{S}) \times H_0^1(\Omega)$ such that

$$(3.34) \quad (\sigma, \tau) + \langle \operatorname{div} \operatorname{div} \tau, u \rangle = 0 \quad \forall \tau \in \mathbf{H}^{-1}(\operatorname{div} \operatorname{div}, \Omega; \mathbb{S}),$$

$$(3.35) \quad \langle \operatorname{div} \operatorname{div} \sigma, v \rangle = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega).$$

We recall the known decoupling from [40, 47], which fits into the framework of this paper as follows. We construct the following commutative diagram from (2.34):

$$\begin{array}{ccc} & & \mathbf{L}^2(\Omega; \mathbb{S}) \\ & & \cup \\ \mathbf{H}^1(\Omega; \mathbb{R}^2) & \xrightarrow{\operatorname{sym} \operatorname{curl}} & \mathbf{H}^{-1}(\operatorname{div} \operatorname{div}, \Omega; \mathbb{S}) \xrightarrow{\operatorname{div} \operatorname{div}} H^{-1}(\Omega) \longrightarrow 0 \quad . \\ & \uparrow \mathbf{r} & \uparrow \Delta \\ & \mathbf{H}_0^1(\Omega; \mathbb{S}) & \xleftarrow{\pi} H_0^1(\Omega) \end{array}$$

According to Theorem 3.4 and Helmholtz decomposition (2.35), the mixed formulation (3.34)–(3.35) can be decoupled to find $w \in H_0^1(\Omega)$, $p \in \mathbf{H}^1(\Omega; \mathbb{R}^2)/\mathbf{RM}^{\operatorname{rot}}$,

and $u \in H_0^1(\Omega)$ such that

$$(3.36) \quad (\nabla w, \nabla v) = -\langle f, v \rangle \quad \forall v \in H_0^1(\Omega),$$

$$(3.37) \quad (\text{sym curl } \mathbf{p}, \text{sym curl } \mathbf{q}) = -(\boldsymbol{\pi} w, \text{sym curl } \mathbf{q}) \quad \forall \mathbf{q} \in \mathbf{H}^1(\Omega; \mathbb{R}^2) / \mathbf{RM}^{\text{rot}},$$

$$(3.38) \quad (\nabla u, \nabla \chi) = (\boldsymbol{\sigma}, \boldsymbol{\pi} \chi) \quad \forall \chi \in H_0^1(\Omega),$$

where $\boldsymbol{\sigma} = \text{sym curl } \mathbf{p} + \boldsymbol{\pi} w$, and

$$\mathbf{RM}^{\text{rot}} := \text{span} \left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \mathbf{x} \right\}.$$

The second equation is also equivalent to the linear elasticity problem

$$(\boldsymbol{\varepsilon}(\mathbf{p}^\perp), \boldsymbol{\varepsilon}(\mathbf{q}^\perp)) = -(\boldsymbol{\pi} w, \boldsymbol{\varepsilon}(\mathbf{q}^\perp)) \quad \forall \mathbf{q}^\perp \in \mathbf{H}^1(\Omega; \mathbb{R}^2) / \mathbf{RM},$$

where rigid motion space

$$\mathbf{RM} := \text{span} \left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \mathbf{x}^\perp \right\}.$$

Such a decomposition was first obtained in [40] and has been recently generalized to the mixed boundary conditions in [47]. Similarly, we can also recover the decomposition of the mixed formulation (3.34)–(3.35) in three dimensions in [46].

3.4. The primal formulation of the fourth order curl equation. Let $\Omega \subset \mathbb{R}^3$ and $\mathbf{f} \in \mathbf{H}(\text{div}, \Omega)$ with $\text{div } \mathbf{f} = 0$. Consider the fourth order curl equation

$$(3.39) \quad \begin{cases} (\text{curl})^4 \mathbf{u} = \mathbf{f} & \text{in } \Omega, \\ \text{div } \mathbf{u} = 0 & \text{in } \Omega, \\ \mathbf{u} \times \boldsymbol{\nu} = (\text{curl } \mathbf{u}) \times \boldsymbol{\nu} = \mathbf{0} & \text{on } \partial\Omega. \end{cases}$$

The primal formulation of (3.39) is to find $\mathbf{u} \in \mathbf{H}_0^2(\text{curl}, \Omega)$ such that

$$(3.40) \quad (\text{curl curl } \mathbf{u}, \text{curl curl } \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in \mathbf{H}_0^2(\text{curl}, \Omega),$$

where

$$\mathbf{H}_0^2(\text{curl}, \Omega) := \{ \mathbf{v} \in \mathbf{L}^2(\Omega, \mathbb{R}^3) : \text{curl } \mathbf{v}, \text{curl curl } \mathbf{v} \in \mathbf{L}^2(\Omega, \mathbb{R}^3), \text{div } \mathbf{v} = 0, \text{ and } \mathbf{v} \times \mathbf{n} = (\text{curl } \mathbf{v}) \times \mathbf{n} = \mathbf{0} \}.$$

Setting up the commutative diagram from (2.30)

$$\begin{array}{ccccccc} \mathbf{H}_0^1(\Omega; \mathbb{R}^3) & \xrightarrow{\Delta} & \mathbf{H}^{-1}(\Omega; \mathbb{R}^3) & & & & \\ & & \cup & & & & \\ L_0^2(\Omega) & \xrightarrow{\text{grad}} & \mathbf{H}^{-1}(\text{curl}, \Omega) & \xrightarrow{\text{curl}} & (K_0^c)' & \longrightarrow & 0 \\ & & \uparrow I & & \uparrow \text{curl curl} & & \\ & & \mathbf{H}_0(\text{div}, \Omega) & \xleftarrow{\text{curl}} & K_0^c & & \end{array}$$

the corresponding mixed formulation is to find $(\boldsymbol{\gamma}, \mathbf{u}) \in \mathbf{H}^{-1}(\text{curl}, \Omega) \times K_0^c$ such that

$$(3.41) \quad (\boldsymbol{\gamma}, \boldsymbol{\beta})_{-1} - \langle \text{curl } \boldsymbol{\beta}, \mathbf{u} \rangle = 0 \quad \forall \boldsymbol{\beta} \in \mathbf{H}^{-1}(\text{curl}, \Omega),$$

$$(3.42) \quad \langle \text{curl } \boldsymbol{\gamma}, \mathbf{v} \rangle = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in K_0^c.$$

By introducing variable ϕ and applying the Helmholtz decomposition (2.31)

$$\phi = -\Delta^{-1}\gamma = -\Delta^{-1}(\operatorname{curl} \mathbf{w} + \nabla p) \in \mathbf{H}_0^1(\Omega; \mathbb{R}^3),$$

from Theorem 3.4 the decoupled and unfolded system is to find $\mathbf{w} \in K_0^c$, $\phi \in \mathbf{H}_0^1(\Omega; \mathbb{R}^3)$, $p \in L_0^2(\Omega)$, and $\mathbf{u} \in K_0^c$ such that

$$(3.43) \quad (\operatorname{curl} \mathbf{w}, \operatorname{curl} \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in K_0^c,$$

$$(3.44) \quad (\nabla \phi, \nabla \psi) + (\operatorname{div} \psi, p) = (\operatorname{curl} \mathbf{w}, \psi) \quad \forall \psi \in \mathbf{H}_0^1(\Omega; \mathbb{R}^3),$$

$$(3.45) \quad (\operatorname{div} \phi, q) = 0 \quad \forall q \in L_0^2(\Omega),$$

$$(3.46) \quad (\operatorname{curl} \mathbf{u}, \operatorname{curl} \chi) = (\phi, \operatorname{curl} \chi) \quad \forall \chi \in K_0^c.$$

According to (3.45)–(3.46), we have $\operatorname{curl} \mathbf{u} = \phi \in \mathbf{H}_0^1(\Omega; \mathbb{R}^3)$. Note that

$$(\nabla \phi, \nabla \psi) = (\operatorname{curl} \phi, \operatorname{curl} \psi) + (\operatorname{div} \phi, \operatorname{div} \psi).$$

Thus we get from (3.44)

$$(\operatorname{curl} \operatorname{curl} \mathbf{u}, \operatorname{curl} \operatorname{curl} \mathbf{v}) = (\operatorname{curl} \mathbf{w}, \operatorname{curl} \mathbf{v})$$

for any $\mathbf{v} \in K_0^c$ satisfying $\operatorname{curl} \mathbf{v} \in \mathbf{H}_0^1(\Omega; \mathbb{R}^3)$. Combined with (3.43), the decoupled formulation (3.43)–(3.46) is equivalent to the primal formulation (3.40) of the fourth order curl equation. The decoupling (3.43)–(3.46) is also presented in [52] independently and is based on a different approach there.

Therefore, we can solve the four curl problem by solving two Maxwell’s equations and one Stokes equation. The Maxwell’s equation with divergence-free constraint can be further decoupled into one vector Poisson equation and one scalar Poisson equation [23].

3.5. The mixed formulation of the fourth order curl equation. A mixed formulation of the fourth order curl equation (3.39) is to find $\phi \in \mathbf{H}(\operatorname{curl} \operatorname{curl}, (K_0^c)')$ and $\mathbf{u} \in K_0^c$ such that

$$(3.47) \quad (\phi, \psi) + \langle \operatorname{curl} \operatorname{curl} \psi, \mathbf{u} \rangle = 0 \quad \forall \psi \in \mathbf{H}(\operatorname{curl} \operatorname{curl}, (K_0^c)'),$$

$$(3.48) \quad \langle \operatorname{curl} \operatorname{curl} \phi, \mathbf{v} \rangle = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in K_0^c.$$

The following commutative diagram is designed from (2.41):

$$\begin{array}{ccccccc} & & & & \mathbf{L}^2(\Omega; \mathbb{R}^3) & & \\ & & & & \cup & & \\ \mathbf{H}^0(\operatorname{div}, \operatorname{dev} \operatorname{sym}) & \xrightarrow{\operatorname{spn}^{-1} \operatorname{skw}} & \mathbf{H}(\operatorname{curl} \operatorname{curl}, (K_0^c)') & \xrightarrow{\operatorname{curl} \operatorname{curl}} & (K_0^c)' & \longrightarrow & 0 \quad . \\ & & \uparrow I & & \uparrow \operatorname{curl} \operatorname{curl} & & \\ & & K_0^c & \xleftarrow{I} & K_0^c & & \end{array}$$

Applying Theorem 3.4 and Helmholtz decomposition (2.42), the mixed formulation (3.47)–(3.48) of the fourth order curl equation is equivalent to finding $\mathbf{w} \in K_0^c$, $\sigma \in \mathbf{H}^0(\operatorname{div}, \operatorname{dev} \operatorname{sym})$, and $\mathbf{u} \in K_0^c$ such that

$$(3.49) \quad (\operatorname{curl} \mathbf{w}, \operatorname{curl} \mathbf{v}) = \langle \mathbf{f}, \mathbf{v} \rangle \quad \forall \mathbf{v} \in K_0^c,$$

$$(3.50) \quad (\operatorname{skw} \sigma, \operatorname{skw} \tau) = -(\operatorname{spn} \mathbf{w}, \operatorname{skw} \tau) \quad \forall \tau \in \mathbf{H}^0(\operatorname{div}, \operatorname{dev} \operatorname{sym}),$$

$$(3.51) \quad (\operatorname{curl} \mathbf{u}, \operatorname{curl} \chi) = -(\phi, \chi) \quad \forall \chi \in K_0^c,$$

where $\phi = \text{spn}^{-1} \text{skw } \sigma + w$.

By (2.40), let $\sigma = \rho I + \text{spn } \alpha$. Then (3.50) is equivalent to finding $\alpha \in L^2(\Omega; \mathbb{R}^3)$, $\rho \in L^2_0(\Omega)$, and $p \in H^1_0(\Omega; \mathbb{R}^3)$ such that

$$\begin{aligned} (\alpha, \beta) + (\gamma I + \text{spn } \beta, \nabla p) &= -(\omega, \beta) & \forall \beta \in L^2(\Omega; \mathbb{R}^3), \gamma \in L^2_0(\Omega), \\ (\rho I + \text{spn } \alpha, \nabla q) &= 0 & \forall q \in H^1_0(\Omega; \mathbb{R}^3). \end{aligned}$$

Again, both (3.49) and (3.51) are Maxwell's equations.

3.6. The primal formulation of the triharmonic equation. Consider the triharmonic equation with homogeneous Dirichlet boundary condition

$$(3.52) \quad \begin{cases} -\Delta^3 u = f & \text{in } \Omega, \\ u = \partial_\nu u = \partial_{\nu\nu} u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $f \in H^{-2}(\Omega)$ with $\Omega \subset \mathbb{R}^2$. The primal formulation of (3.52) is to find $u \in H^3_0(\Omega)$ such that

$$(3.53) \quad (\nabla^3 u, \nabla^3 v) = \langle f, v \rangle \quad \forall v \in H^3_0(\Omega).$$

In view of the following commutative diagram derived from (2.37),

$$\begin{array}{ccccccc} H^1_0(\Omega; \mathbb{S}) & \xrightarrow{\Delta} & H^{-1}(\Omega; \mathbb{S}) & & & & \\ & & \cup & & & & \\ L^2(\Omega; \mathbb{R}^2) & \xrightarrow{\varepsilon} & H^{-2}(\text{rot rot}, \Omega; \mathbb{S}) & \xrightarrow{\text{rot rot}} & H^{-2}(\Omega) & \longrightarrow & 0 \\ & & \uparrow I & & \uparrow \Delta^2 & & \\ & & H_0(\text{div}, \Omega; \mathbb{S}) & \xleftarrow{\text{curl curl}} & H^2_0(\Omega) & & \end{array}$$

the corresponding mixed formulation is to find $(\gamma, u) \in H^{-2}(\text{rot rot}, \Omega; \mathbb{S}) \times H^2_0(\Omega)$ such that

$$(3.54) \quad (\gamma, \beta)_{-1} - \langle \text{rot rot } \beta, u \rangle = 0 \quad \forall \beta \in H^{-2}(\text{rot rot}, \Omega; \mathbb{S}),$$

$$(3.55) \quad \langle \text{rot rot } \gamma, v \rangle = \langle f, v \rangle \quad \forall v \in H^2_0(\Omega).$$

By introducing a variable and applying the Helmholtz decomposition (2.38),

$$\phi = -\Delta^{-1} \gamma = -\Delta^{-1} (\varepsilon(p) + \text{curl curl } w) \in H^1_0(\Omega; \mathbb{S}),$$

from Theorem 3.4, the decoupled system is to find $w \in H^2_0(\Omega)$, $\phi \in H^1_0(\Omega; \mathbb{S})$, $p \in L^2(\Omega; \mathbb{R}^2)/\mathbf{RM}$, and $u \in H^2_0(\Omega)$ such that

$$(3.56) \quad (\text{curl curl } w, \text{curl curl } v) = \langle f, v \rangle \quad \forall v \in H^2_0(\Omega),$$

$$(3.57) \quad (\nabla \phi, \nabla \psi) + (\text{div } \psi, p) = (\text{curl curl } w, \psi) \quad \forall \psi \in H^1_0(\Omega; \mathbb{S}),$$

$$(3.58) \quad (\text{div } \phi, q) = 0 \quad \forall q \in L^2(\Omega; \mathbb{R}^2)/\mathbf{RM},$$

$$(3.59) \quad (\text{curl curl } u, \text{curl curl } \chi) = (\phi, \text{curl curl } \chi) \quad \forall \chi \in H^2_0(\Omega).$$

It is evident that the primal formulation (3.53) of the triharmonic equation is equivalent to two biharmonic equations and one Stokes equation (cf. (3.56)–(3.59)), which is different from the decoupling in [27]. These equations can be discretized directly, since there exist many finite elements for discretizing the biharmonic equations in the literature.

Recursively applying the decomposition, we can decouple the m th harmonic equation $\Delta^m u = f$ with homogeneous Dirichlet boundary condition, i.e., $u \in H^m_0(\Omega)$, into a sequence of Poisson and Stokes equations.

3.7. The mixed formulation of the triharmonic equation. A mixed formulation of the triharmonic equation (3.52) in two dimensions is to find $(\boldsymbol{\sigma}, u) \in \mathbf{H}^{-2}(\text{div}^3, \Omega) \times H_0^2(\Omega)$ such that

$$(3.60) \quad (\boldsymbol{\sigma}, \boldsymbol{\tau}) + \langle \text{div div div } \boldsymbol{\tau}, u \rangle = 0 \quad \forall \boldsymbol{\tau} \in \mathbf{H}^{-2}(\text{div}^3, \Omega),$$

$$(3.61) \quad \langle \text{div div div } \boldsymbol{\sigma}, v \rangle = \langle f, v \rangle \quad \forall v \in H_0^2(\Omega).$$

Due to (2.43), we construct the commutative diagram

$$\begin{array}{ccccccc} & & & L^2(\Omega; \mathbb{S}(3)) & & & \\ & & & \cup & & & \\ \mathbf{H}^1(\Omega; \mathbb{S}) & \xrightarrow{\text{sym curl}} & \mathbf{H}^{-2}(\text{div}^3, \Omega) & \xrightarrow{\text{div div div}} & H^{-2}(\Omega) & \longrightarrow & 0 \quad . \\ & & \Xi \uparrow & & \uparrow \Delta^2 & & \\ & & H_0^1(\Omega; \mathbb{R}^2) & \xleftarrow{\nabla} & H_0^2(\Omega) & & \end{array}$$

Then, according to Theorem 3.4 and the Helmholtz decomposition (2.44), the mixed formulation (3.60)–(3.61) of the triharmonic equation is equivalent to finding $w \in H_0^2(\Omega)$, $\boldsymbol{\phi} \in \mathbf{H}^1(\Omega; \mathbb{S})$, and $u \in H_0^2(\Omega)$ such that

$$(3.62) \quad (\Delta w, \Delta v) = \langle f, v \rangle \quad \forall v \in H_0^2(\Omega),$$

$$(3.63) \quad (\text{sym curl } \boldsymbol{\phi}, \text{sym curl } \boldsymbol{\psi}) = -(\Xi \nabla w, \text{sym curl } \boldsymbol{\psi}) \quad \forall \boldsymbol{\psi} \in \mathbf{H}^1(\Omega; \mathbb{S}),$$

$$(3.64) \quad (\Delta u, \Delta \chi) = -(\boldsymbol{\sigma}, \Xi \nabla \chi) \quad \forall \chi \in H_0^2(\Omega),$$

where $\boldsymbol{\sigma} = \text{sym curl } \boldsymbol{\phi} + \Xi \nabla w$.

4. Discretization based on decoupled formulation. In this section, we will consider discretization based on the decoupled formulation. By decoupling the fourth order equation into second order equations, we can use well-known conforming finite element spaces. Furthermore, we can easily derive the superconvergence to the Galerkin projections without any mesh conditions, which is not known in the literature.

4.1. Decoupled discretization of HHJ formulation. Let $f \in L^2(\Omega)$, $V_h \subset H_0^1(\Omega)$, and $\mathbf{P}_h \subset \mathbf{H}^1(\Omega; \mathbb{R}^2)$. The discrete method based on formulation (3.36)–(3.38) is to find $w_h \in V_h$, $\mathbf{p}_h \in \mathbf{P}_h/\mathbf{RM}^{\text{rot}}$, and $u_h \in V_h$ such that

$$(4.1) \quad (\nabla w_h, \nabla v_h) = -(f, v_h) \quad \forall v_h \in V_h,$$

$$(4.2) \quad (\text{sym curl } \mathbf{p}_h, \text{sym curl } \mathbf{q}_h) = -(\boldsymbol{\pi} w_h, \text{sym curl } \mathbf{q}_h) \quad \forall \mathbf{q}_h \in \mathbf{P}_h/\mathbf{RM}^{\text{rot}},$$

$$(4.3) \quad (\nabla u_h, \nabla \chi_h) = (\boldsymbol{\sigma}_h, \boldsymbol{\pi} \chi_h) \quad \forall \chi_h \in V_h,$$

where $\boldsymbol{\sigma}_h = \text{sym curl } \mathbf{p}_h + \boldsymbol{\pi} w_h$.

Define projection $\mathbf{P}_h^{cs} : \mathbf{H}^1(\Omega; \mathbb{R}^2) \rightarrow \mathbf{P}_h/\mathbf{RM}^{\text{rot}}$ by

$$(\text{sym curl } \mathbf{P}_h^{cs} \mathbf{p}, \text{sym curl } \mathbf{q}_h) = (\text{sym curl } \mathbf{p}, \text{sym curl } \mathbf{q}_h).$$

Similarly, denote by P_h^{grad} the H^1 orthogonal projection onto V_h . Let

$$\boldsymbol{\sigma}_h^* := \text{sym curl } \mathbf{P}_h^{cs} \mathbf{p} + \boldsymbol{\pi} w_h.$$

LEMMA 4.1. *Let (w, \mathbf{p}, u) be the solution of HHJ mixed formulation (3.36)–(3.38) and (w_h, \mathbf{p}_h, u_h) be the solution of (4.1)–(4.3). We then have the estimates*

$$|w - w_h|_1 \lesssim \inf_{v_h \in V_h} |w - v_h|_1,$$

$$\|\text{sym curl}(\mathbf{P}_h^{cs} \mathbf{p} - \mathbf{p}_h)\|_0 + \|\boldsymbol{\sigma}_h^* - \boldsymbol{\sigma}_h\|_0 \lesssim \|w - w_h\|_0,$$

$$|P_h^{\text{grad}} u - u_h|_1 \lesssim \|\mathbf{p} - \mathbf{p}_h\|_0 + \|w - w_h\|_0.$$

Proof. Subtracting (4.1)–(4.3) from (3.36)–(3.38), we get the error equations

$$\begin{aligned} (\nabla(w - w_h), \nabla v_h) &= 0 & \forall v_h \in V_h, \\ (\text{sym curl}(\mathbf{P}_h^{cs} \mathbf{p} - \mathbf{p}_h), \text{sym curl} \mathbf{q}_h) &= (\boldsymbol{\pi}(w_h - w), \text{sym curl} \mathbf{q}_h) & \forall \mathbf{q}_h \in \mathbf{P}_h / \mathbf{RM}^{\text{rot}}, \\ (\nabla(P_h^{\text{grad}} u - u_h), \nabla \chi_h) &= (\boldsymbol{\sigma} - \boldsymbol{\sigma}_h, \boldsymbol{\pi} \chi_h) & \forall \chi_h \in V_h. \end{aligned}$$

Then all the error estimates hold by standard argument. □

Furthermore, assume

$$(4.4) \quad \|w - w_h\|_0 \lesssim h^\delta |w - w_h|_1, \quad \|\mathbf{p} - \mathbf{p}_h\|_0 \lesssim h^\delta \|\text{sym curl}(\mathbf{p} - \mathbf{p}_h)\|_0,$$

where $\delta \in (1/2, 1]$ is the regularity constant depending on the shape of Ω . This assumption can be proved by the duality argument (cf. [24, 10]).

THEOREM 4.2. *Let (w, \mathbf{p}, u) be the solution of HHJ mixed formulation (3.36)–(3.38) and (w_h, \mathbf{p}_h, u_h) be the solution of (4.1)–(4.3). We then have the estimates*

$$\|\text{sym curl}(\mathbf{p} - \mathbf{p}_h)\|_0 + \|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\|_0 \lesssim \inf_{\mathbf{q}_h \in \mathbf{P}_h} \|\text{sym curl}(\mathbf{p} - \mathbf{q}_h)\|_0 + \inf_{v_h \in V_h} |w - v_h|_1,$$

$$|u - u_h|_1 \lesssim \inf_{v_h \in V_h} |u - v_h|_1 + \inf_{\mathbf{q}_h \in \mathbf{P}_h} \|\text{sym curl}(\mathbf{p} - \mathbf{q}_h)\|_0 + \inf_{v_h \in V_h} |w - v_h|_1.$$

Moreover if assumption (4.4) is true, we have the improved error estimates

$$(4.5) \quad \|\text{sym curl}(\mathbf{P}_h^{cs} \mathbf{p} - \mathbf{p}_h)\|_0 + \|\boldsymbol{\sigma}_h^* - \boldsymbol{\sigma}_h\|_0 \lesssim h^\delta \inf_{v_h \in V_h} |w - v_h|_1,$$

$$(4.6) \quad |P_h^{\text{grad}} u - u_h|_1 \lesssim h^\delta \inf_{\mathbf{q}_h \in \mathbf{P}_h} \|\text{sym curl}(\mathbf{p} - \mathbf{q}_h)\|_0 + h^\delta \inf_{v_h \in V_h} |w - v_h|_1.$$

Proof. The first two error estimates can be derived from Lemma 4.1 and the Poincaré inequality. We can acquire (4.5)–(4.6) from Lemma 4.1 and (4.4). □

Remark 4.3. The error estimates (4.5)–(4.6) are superconvergent if we use equal order finite element spaces for V_h and \mathbf{P}_h .

4.2. Decoupled discretization for biharmonic equation. Now we discretize formulation (3.29)–(3.32) using more general finite element spaces.

Let $f \in L^2(\Omega)$, $V_h \subset H_0^1(\Omega)$, $\mathbf{X}_h \subset \mathbf{H}_0^1(\Omega; \mathbb{R}^2)$, and $P_h \subset L_0^2(\Omega)$. The discrete method based on formulation (3.29)–(3.32) is to find $w_h, u_h \in V_h$, $\boldsymbol{\phi}_h \in \mathbf{X}_h$, and $p_h \in P_h$ such that

$$(4.7) \quad (\text{curl } w_h, \text{curl } v_h) = (f, v_h) \quad \forall v_h \in V_h.$$

$$(4.8) \quad (\nabla \boldsymbol{\phi}_h, \nabla \boldsymbol{\psi}_h) + (\text{div } \boldsymbol{\psi}_h, p_h) = (\text{curl } w_h, \boldsymbol{\psi}_h) \quad \forall \boldsymbol{\psi}_h \in \mathbf{X}_h,$$

$$(4.9) \quad (\text{div } \boldsymbol{\phi}_h, q_h) = 0 \quad \forall q_h \in P_h,$$

$$(4.10) \quad (\text{curl } u_h, \text{curl } \chi_h) = (\boldsymbol{\phi}_h, \text{curl } \chi_h) \quad \forall \chi_h \in V_h.$$

We assume (\mathbf{X}_h, P_h) is a stable finite element pair for the Stokes equation (cf. [8, 7]); i.e., the following inf-sup condition holds:

$$(4.11) \quad \|q_h\|_0 \lesssim \sup_{\boldsymbol{\psi}_h \in \boldsymbol{\Sigma}_h} \frac{(\operatorname{div} \boldsymbol{\psi}_h, q_h)}{|\boldsymbol{\psi}_h|_1} \quad \forall q_h \in P_h.$$

To analyze the discrete method (4.7)–(4.10), we rewrite it as a mixed finite element method

$$\begin{aligned} a(\boldsymbol{\phi}_h, u_h; \boldsymbol{\psi}_h, v_h) + b(\boldsymbol{\psi}_h, v_h; p_h, w_h) &= (f, v_h) \quad \forall (\boldsymbol{\psi}_h, v_h) \in \mathbf{X}_h \times V_h, \\ b(\boldsymbol{\phi}_h, u_h; q_h, \chi_h) &= 0 \quad \forall (q_h, \chi_h) \in P_h \times V_h, \end{aligned}$$

where

$$\begin{aligned} a(\boldsymbol{\phi}_h, u_h; \boldsymbol{\psi}_h, v_h) &:= (\nabla \boldsymbol{\phi}_h, \nabla \boldsymbol{\psi}_h), \\ b(\boldsymbol{\phi}_h, u_h; q_h, \chi_h) &:= (\operatorname{div} \boldsymbol{\phi}_h, q_h) - (\boldsymbol{\phi}_h, \operatorname{curl} \chi_h) + (\operatorname{curl} u_h, \operatorname{curl} \chi_h). \end{aligned}$$

LEMMA 4.4. *Assume the inf-sup condition (4.11); the following inf-sup condition holds:*

$$(4.12) \quad \|q_h\|_0 + |\chi_h|_1 \lesssim \sup_{(\boldsymbol{\psi}_h, v_h) \in \boldsymbol{\Sigma}_h \times V_h} \frac{b(\boldsymbol{\psi}_h, v_h; q_h, \chi_h)}{|\boldsymbol{\psi}_h|_1 + |v_h|_1} \quad \forall (q_h, \chi_h) \in P_h \times V_h.$$

Proof. It is easy to see that

$$|\chi_h|_1 = \sup_{v_h \in V_h} \frac{(\operatorname{curl} v_h, \operatorname{curl} \chi_h)}{|v_h|_1} \leq \sup_{(\boldsymbol{\psi}_h, v_h) \in \boldsymbol{\Sigma}_h \times V_h} \frac{b(\boldsymbol{\psi}_h, v_h; q_h, \chi_h)}{|\boldsymbol{\psi}_h|_1 + |v_h|_1}.$$

It follows from (4.11) and the Poincaré inequality that

$$\begin{aligned} \|q_h\|_0 &\lesssim \sup_{\boldsymbol{\psi}_h \in \boldsymbol{\Sigma}_h} \frac{(\operatorname{div} \boldsymbol{\psi}_h, q_h)}{|\boldsymbol{\psi}_h|_1} = \sup_{\boldsymbol{\psi}_h \in \boldsymbol{\Sigma}_h} \frac{b(\boldsymbol{\psi}_h, 0; q_h, \chi_h) + (\boldsymbol{\psi}_h, \operatorname{curl} \chi_h)}{|\boldsymbol{\psi}_h|_1} \\ &\lesssim |\chi_h|_1 + \sup_{(\boldsymbol{\psi}_h, v_h) \in \boldsymbol{\Sigma}_h \times V_h} \frac{b(\boldsymbol{\psi}_h, v_h; q_h, \chi_h)}{|\boldsymbol{\psi}_h|_1 + |v_h|_1}. \end{aligned}$$

Therefore, the inf-sup condition (4.12) will be derived by combining the last two inequalities. \square

THEOREM 4.5. *Let $(w, \boldsymbol{\phi}, p, u)$ be the solution of the mixed formulation (3.29)–(3.32), and let $(w_h, \boldsymbol{\phi}_h, p_h, u_h) \in V_h \times \mathbf{X}_h \times P_h \times V_h$ be the solution of the discrete method (4.7)–(4.10). Assume both V_h and \mathbf{X}_h are H^1 conforming, the inf-sup condition (4.11) holds, and the discrete spaces are consistent with respect to the mixed formulation (3.29)–(3.32); then*

$$\begin{aligned} (4.13) \quad &\|w - w_h\|_1 + \|\boldsymbol{\phi} - \boldsymbol{\phi}_h\|_1 + \|p - p_h\|_0 + \|u - u_h\|_1 \\ (4.14) \quad &\lesssim \inf_{\chi_h \in V_h} \|w - \chi_h\|_1 + \inf_{\boldsymbol{\psi}_h \in \mathbf{X}_h} \|\boldsymbol{\phi} - \boldsymbol{\psi}_h\|_1 + \inf_{q_h \in P_h} \|p - q_h\|_0 + \inf_{v_h \in V_h} \|u - v_h\|_1. \end{aligned}$$

Moreover, if

$$(4.15) \quad \|\boldsymbol{\phi} - \boldsymbol{\phi}_h\|_0 \lesssim h^\delta \left(\|\boldsymbol{\phi} - \boldsymbol{\phi}_h\|_1 + \inf_{q_h \in P_h} \|p - q_h\|_0 \right),$$

then

$$(4.16) \quad |P_h^{\operatorname{grad}} u - u_h|_1 \lesssim h^\delta \left(\inf_{\chi_h \in V_h} \|w - \chi_h\|_1 + \inf_{\boldsymbol{\psi}_h \in \mathbf{X}_h} \|\boldsymbol{\phi} - \boldsymbol{\psi}_h\|_1 + \inf_{q_h \in P_h} \|p - q_h\|_0 \right).$$

Proof. For any $(\boldsymbol{\psi}_h, v_h) \in \Sigma_h \times V_h$ satisfying

$$b(\boldsymbol{\psi}_h, v_h; q_h, \chi_h) = 0 \quad \forall (q_h, \chi_h) \in P_h \times V_h,$$

we have

$$(\boldsymbol{\psi}_h, \operatorname{curl} \chi_h) = (\operatorname{curl} v_h, \operatorname{curl} \chi_h) \quad \forall \chi_h \in V_h,$$

which implies

$$|v_h|_1 \leq \|\boldsymbol{\psi}_h\|_0 \lesssim |\boldsymbol{\psi}_h|_1.$$

Thus

$$|\boldsymbol{\psi}_h|_1^2 + |v_h|_1^2 \lesssim |\boldsymbol{\psi}_h|_1^2 = a(\boldsymbol{\psi}_h, v_h; \boldsymbol{\psi}_h, v_h).$$

Combining the above with the inf-sup condition (4.12), we will obtain the error estimate (4.13) by the mixed finite element method theory in [8], and (4.16) can be derived using the similar argument adopted in section 4.1. \square

REFERENCES

- [1] A. ALONSO, *Error estimators for a mixed method*, Numer. Math., 74 (1996), pp. 385–395, <https://doi.org/10.1007/s002110050222>.
- [2] C. AMROUCHE, P. G. CIARLET, AND P. CIARLET, JR., *Weak vector and scalar potentials: Applications to Poincaré’s theorem and Korn’s inequality in Sobolev spaces with negative exponents*, Anal. Appl. (Singap.), 8 (2010), pp. 1–17, <https://doi.org/10.1142/S0219530510001497>.
- [3] D. N. ARNOLD, R. S. FALK, AND R. WINTHER, *Multigrid in $H(\operatorname{div})$ and $H(\operatorname{curl})$* , Numer. Math., 85 (2000), pp. 197–217, <https://doi.org/10.1007/PL00005386>.
- [4] D. N. ARNOLD, R. S. FALK, AND R. WINTHER, *Finite element exterior calculus, homological techniques, and applications*, Acta Numer., 15 (2006), pp. 1–155, <https://doi.org/10.1017/S0962492906210018>.
- [5] D. N. ARNOLD, R. S. FALK, AND R. WINTHER, *Finite element exterior calculus: From Hodge theory to numerical stability*, Bull. Amer. Math. Soc. (N.S.), 47 (2010), pp. 281–354, <https://doi.org/10.1090/S0273-0979-10-01278-4>.
- [6] I. BABUŠKA AND A. K. AZIZ, *Survey lectures on the mathematical foundations of the finite element method*, in The Mathematical Foundations of the Finite Element Method with Applications to Partial Differential Equations (Univ. Maryland, Baltimore, MD, 1972), Academic Press, New York, 1972, pp. 1–359.
- [7] D. BOFFI, F. BREZZI, L. F. DEMKOWICZ, R. G. DURÁN, R. S. FALK, AND M. FORTIN, *Mixed Finite Elements, Compatibility Conditions, and Applications*, Lecture Notes in Math. 1939, Springer-Verlag, Berlin, 2008, <https://doi.org/10.1007/978-3-540-78319-0>.
- [8] D. BOFFI, F. BREZZI, AND M. FORTIN, *Mixed Finite Element Methods and Applications*, Springer, Heidelberg, 2013, <https://doi.org/10.1007/978-3-642-36519-5>.
- [9] D. BRAESS, *Finite Elements: Theory, Fast Solvers, and Applications in Elasticity Theory*, 3rd ed., Cambridge University Press, Cambridge, UK, 2007, <https://doi.org/10.1017/CBO9780511618635>.
- [10] S. C. BRENNER AND L. R. SCOTT, *The Mathematical Theory of Finite Element Methods*, 3rd ed., Springer, New York, 2008, <https://doi.org/10.1007/978-0-387-75934-0>.
- [11] S. C. BRENNER, J. SUN, AND L.-Y. SUNG, *Hodge decomposition methods for a quad-curl problem on planar domains*, J. Sci. Comput., 73 (2017), pp. 495–513, <https://doi.org/10.1007/s10915-017-0449-0>.
- [12] H. BREZIS, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Universitext, Springer, New York, 2011, <https://doi.org/10.1007/978-0-387-70914-7>.
- [13] F. BREZZI, *On the existence, uniqueness and approximation of saddle-point problems arising from Lagrangian multipliers*, Rev. Française Automat. Informat. Recherche Opérationnelle Sér. Rouge, 8 (1974), pp. 129–151.
- [14] F. BREZZI AND M. FORTIN, *Numerical approximation of Mindlin-Reissner plates*, Math. Comp., 47 (1986), pp. 151–158, <https://doi.org/10.2307/2008086>.

- [15] F. BREZZI, M. FORTIN, AND R. STENBERG, *Error analysis of mixed-interpolated elements for Reissner-Mindlin plates*, Math. Models Methods Appl. Sci., 1 (1991), pp. 125–151, <https://doi.org/10.1142/S0218202591000083>.
- [16] C. CARSTENSEN, *A unifying theory of a posteriori finite element error control*, Numer. Math., 100 (2005), pp. 617–637, <https://doi.org/10.1007/s00211-004-0577-y>.
- [17] C. CARSTENSEN AND S. BARTELS, *Each averaging technique yields reliable a posteriori error control in FEM on unstructured grids. I. Low order conforming, nonconforming, and mixed FEM*, Math. Comp., 71 (2002), pp. 945–969, <https://doi.org/10.1090/S0025-5718-02-01402-3>.
- [18] C. CARSTENSEN, S. BARTELS, AND S. JANSCHKE, *A posteriori error estimates for nonconforming finite element methods*, Numer. Math., 92 (2002), pp. 233–256, <https://doi.org/10.1007/s002110100378>.
- [19] C. CARSTENSEN AND J. HU, *A unifying theory of a posteriori error control for nonconforming finite element methods*, Numer. Math., 107 (2007), pp. 473–502, <https://doi.org/10.1007/s00211-007-0068-z>.
- [20] L. CHEN, *Multigrid Methods for Constrained Minimization Problems and Application to Saddle Point Problems*, preprint, <https://arxiv.org/abs/1601.04091>, 2016.
- [21] L. CHEN, J. HU, AND X. HUANG, *Multigrid methods for Hellan–Herrmann–Johnson mixed method of Kirchhoff plate bending problems*, J. Sci. Comput., 76 (2018), pp. 673–696, <https://doi.org/10.1007/s10915-017-0636-z>.
- [22] L. CHEN, J. HU, X. HUANG, AND H. MAN, *Residual-based a posteriori error estimates for symmetric conforming mixed finite elements for linear elasticity problems*, Sci. China Math., 61 (2018), pp. 973–992, <https://doi.org/10.1007/s11425-017-9181-2>.
- [23] L. CHEN, Y. WU, L. ZHONG, AND J. ZHOU, *Multigrid preconditioners for mixed finite element methods of vector Laplacian*, J. Sci. Comput., (2018), pp. 1–28, <https://doi.org/10.1007/s10915-018-0697-7>.
- [24] P. G. CIARLET, *The Finite Element Method for Elliptic Problems*, North–Holland, Amsterdam, 1978; Classics Appl. Math. 40, SIAM, Philadelphia, 2002, <https://doi.org/10.1137/1.9780898719208>.
- [25] M. COSTABEL AND A. MCINTOSH, *On Bogovskii and regularized Poincaré integral operators for de Rham complexes on Lipschitz domains*, Math. Z., 265 (2010), pp. 297–320, <https://doi.org/10.1007/s00209-009-0517-8>.
- [26] R. DE LA MBYADRID MODINO, *Quantum Mechanics in Rigged Hilbert Space Language*, Ph.D. thesis, Universidad de Valladolid, Valladolid, Spain, 2001.
- [27] D. GALLISTL, *Stable splitting of polyharmonic operators by generalized Stokes systems*, Math. Comp., 86 (2017), pp. 2555–2577, <https://doi.org/10.1090/mcom/3208>.
- [28] D. GALLISTL, *Variational formulation and numerical analysis of linear elliptic equations in nondivergence form with Cordes coefficients*, SIAM J. Numer. Anal., 55 (2017), pp. 737–757, <https://doi.org/10.1137/16M1080495>.
- [29] V. GIRAULT AND P.-A. RAVIART, *Finite Element Methods for Navier-Stokes Equations: Theory and Algorithms*, Springer-Verlag, Berlin, 1986, <https://doi.org/10.1007/978-3-642-61623-5>.
- [30] A. HATCHER, *Algebraic Topology*, Cambridge University Press, Cambridge, UK, 2002.
- [31] K. HELLAN, *Analysis of Elastic Plates in Flexure by a Simplified Finite Element Method*, Acta Polytechnica Scandinavica Civil Engrg. Building Construct. Ser. 46, Norges tekniske vitenskapsakademi, Trondheim, Norway, 1967.
- [32] L. R. HERRMANN, *Finite element bending analysis for plates*, J. Engrg. Mech. Div., 93 (1967), pp. 49–83.
- [33] R. HIPTMAIR, *Multigrid method for $\mathbf{H}(\text{div})$ in three dimensions*, Electron. Trans. Numer. Anal., 6 (1997), pp. 133–152, <http://etna.mcs.kent.edu/vol.6.1997/pp133-152.dir/pp133-152.html>.
- [34] R. HIPTMAIR, *Finite elements in computational electromagnetism*, Acta Numer., 11 (2002), pp. 237–339, <https://doi.org/10.1017/S0962492902000041>.
- [35] R. HIPTMAIR AND J. XU, *Nodal auxiliary space preconditioning in $\mathbf{H}(\text{curl})$ and $\mathbf{H}(\text{div})$ spaces*, SIAM J. Numer. Anal., 45 (2007), pp. 2483–2509, <https://doi.org/10.1137/060660588>.
- [36] J. HUANG, X. HUANG, AND J. XU, *An Efficient Poisson-Based Solver for Biharmonic Equations Discretized by the Morley Element Method*, Tech. report, 2012.
- [37] J. HUANG, X. HUANG, AND Y. XU, *Convergence of an adaptive mixed finite element method for Kirchhoff plate bending problems*, SIAM J. Numer. Anal., 49 (2011), pp. 574–607, <https://doi.org/10.1137/090773374>.
- [38] X. HUANG, *New Finite Element Methods and Efficient Algorithms for Fourth Order Elliptic Equations*, Ph.D. thesis, Shanghai Jiao Tong University, Shanghai Shi, China, 2010.

- [39] C. JOHNSON, *On the convergence of a mixed finite-element method for plate bending problems*, Numer. Math., 21 (1973), pp. 43–62, <https://doi.org/10.1007/BF01436186>.
- [40] W. KRENDL, K. RAFETSEDER, AND W. ZULEHNER, *A decomposition result for biharmonic problems and the Hellan-Herrmann-Johnson method*, Electron. Trans. Numer. Anal., 45 (2016), pp. 257–282, <http://etna.math.kent.edu/volumes/2011-2020/vol45/abstract.php?vol=45&pages=257-282>.
- [41] K.-A. MARDAL, J. SCHÖBERL, AND R. WINTHER, *A Uniform inf-sup Condition with Applications to Preconditioning*, preprint, <https://arxiv.org/abs/1201.1513>, 2012.
- [42] K. A. MARDAL, X.-C. TAI, AND R. WINTHER, *A robust finite element method for Darcy-Stokes flow*, SIAM J. Numer. Anal., 40 (2002), pp. 1605–1631, <https://doi.org/10.1137/S0036142901383910>.
- [43] K.-A. MARDAL AND R. WINTHER, *Uniform preconditioners for the time dependent Stokes problem*, Numer. Math., 98 (2004), pp. 305–327, <https://doi.org/10.1007/s00211-004-0529-6>.
- [44] P. MONK, *Finite Element Methods for Maxwell's Equations*, Oxford University Press, New York, 2003, <https://doi.org/10.1093/acprof:oso/9780198508885.001.0001>.
- [45] M. A. OLSHANSKII, J. PETERS, AND A. REUSKEN, *Uniform preconditioners for a parameter dependent saddle point problem with application to generalized Stokes interface equations*, Numer. Math., 105 (2006), pp. 159–191, <https://doi.org/10.1007/s00211-006-0031-4>.
- [46] D. PAULY AND W. ZULEHNER, *On Closed and Exact Grad-grad- and div-Div-Complexes, Corresponding Compact Embeddings for Tensor Rotations, and a Related Decomposition Result for Biharmonic Problems in 3D*, <https://arxiv.org/abs/1609.05873>, 2016.
- [47] K. RAFETSEDER AND W. ZULEHNER, *A decomposition result for Kirchhoff plate bending problems and a new discretization approach*, SIAM J. Numer. Anal., 56 (2018), pp. 1961–1986, <https://doi.org/10.1137/17M1118427>.
- [48] M. SCHEDENSACK, *A new discretization for m th-Laplace equations with arbitrary polynomial degrees*, SIAM J. Numer. Anal., 54 (2016), pp. 2138–2162, <https://doi.org/10.1137/15M1013651>.
- [49] H. VON HELMHOLTZ, *On integrals of the hydrodynamic equations that correspond to vortex motions*, Journal für die reine und angewandte Mathematik, 55 (1858), pp. 25–55.
- [50] X. XIE, J. XU, AND G. XUE, *Uniformly-stable finite element methods for Darcy-Stokes-Brinkman models*, J. Comput. Math., 26 (2008), pp. 437–455.
- [51] K. YOSIDA, *Functional Analysis*, 6th ed., Springer-Verlag, Berlin, 1980.
- [52] S. ZHANG, *Mixed schemes for quad-curl equations*, ESAIM Math. Model. Numer. Anal., 52 (2018), pp. 147–161, <https://doi.org/10.1051/m2an/2018005>.
- [53] S. ZHANG, *Regular decomposition and a framework of order reduced methods for fourth order problems*, Numer. Math., 138 (2018), pp. 241–271, <https://doi.org/10.1007/s00211-017-0902-x>.
- [54] S. ZHANG, Y. XI, AND X. JI, *A multi-level mixed element method for the eigenvalue problem of biharmonic equation*, J. Sci. Comput., 75 (2018), pp. 1415–1444, <https://doi.org/10.1007/s10915-017-0592-7>.