

### Assignment 3

---

1. Let  $C^\infty[-1, 1] \ni a \geq a_0 > 0$  and  $y \in [-1, 1]$ . Compute  $G(\cdot, y)$  satisfying

$$-\partial_x(a(x)\partial_x G(\cdot, y)) = \delta_y$$

and  $G(-1, y) = G(1, y) = 0$ . Let  $f \in L^1(-1, 1)$  and show that  $u$  defined through

$$u(x) = \int_{-1}^1 G(x, y)f(y) dy, \quad x \in (-1, 1)$$

solves (in which sense?) the following boundary value problem

$$\begin{cases} -\partial_x(a(x)\partial_x u) = f(x), & x \in (-1, 1), \\ u(-1) = u(1) = 0. \end{cases}$$

2. Let  $\omega \in \mathbb{R}$  and find the solution of

$$(\partial_x^2 + \omega^2)G = \delta \text{ in } \mathcal{D}'(\mathbb{R})$$

[Hint: Let  $G = Hf$  for  $f \in C^\infty(\mathbb{R})$  and the Heaviside function  $H$ ]

3. Let  $T \in \mathcal{E}'(\mathbb{R}^n)$  with  $\text{supp}(T) = K$  be of finite order  $m \in \mathbb{N}$ . Show that

$$\langle T, \phi \rangle = 0 \text{ for any } \phi \in \mathcal{E}(\mathbb{R}^n) \text{ with } \partial^\alpha \phi|_K \equiv 0 \text{ for } |\alpha| \leq m.$$

4. Let  $E$  be a given vector space. A function  $p : E \rightarrow [0, \infty)$  is called *seminorm* if

$$p(x + y) \leq p(x) + p(y) \text{ and } p(\alpha x) = |\alpha|p(x)$$

for  $x, y \in E$  and  $\alpha \in \mathbb{K}$ . A family of seminorms  $\{p_\lambda : \lambda \in \Lambda\}$  is called *separating* if

$$p_\lambda(x) = 0 \text{ for each } \lambda \in \Lambda \text{ implies } x = 0.$$

Now, a set  $X \subset E$  is called *open* if for each  $x \in X$  there are finitely many  $\lambda_j \in \Lambda$  and  $\epsilon_j > 0$  such that

$$x + \bigcap_{j=1, \dots, m} p_{\lambda_j}^{-1}([0, \epsilon_j]) \subset X.$$

Show that the collection of all open sets is a topology on  $E$ .  $E$  endowed with this topology is called *locally convex space*. A linear map

$$u : E \rightarrow E$$

is then continuous iff preimages of open sets are open. Find a characterization of continuity by means of inequalities for the seminorms

$p_\lambda$ . Prove that the space  $\mathcal{E}(\Omega)$  defined in class is a locally convex space with seminorms

$$\{p_{K,m} : K = \bar{K} \subset\subset \Omega, m \in \mathbb{N}\}$$

defined through  $p_{K,m}(\varphi) = \sup_{x \in K, |\alpha| \leq m} |D^\alpha \varphi(x)|$ .

5. Let  $\psi \in \mathcal{D} = \mathcal{D}(\mathbb{R}^n)$  with  $\psi = 1$  on  $\mathbb{B}(0,1)$  and define  $\psi_k$  through

$$\psi_k(x) = \psi\left(\frac{x}{k}\right), x \in \mathbb{R}^n.$$

Show that  $\psi_k \rightarrow \mathbf{1}$  ( $k \rightarrow \infty$ ) in  $\mathcal{E} = \mathcal{E}(\mathbb{R}^n)$ . Then prove that  $\mathcal{D}$  is dense in  $\mathcal{E}$  as well as  $\mathcal{E}'$  in  $\mathcal{D}'$ .