

## Assignment 13

---

1. Assume that  $f \in C([0, 1])$  and that

$$F \in C([0, 1] \times [0, 1] \times \mathbb{R}), \quad \partial_u F \in C([0, 1] \times [0, 1] \times \mathbb{R})$$

and consider the integral equation

$$u(x) = \int_0^1 F(x, y, u(y)) dy + f(x), \quad 0 \leq x \leq 1.$$

Show that, if  $\|\partial_u F\|_\infty < 1$ , the integral equation has a unique solution  $u \in C([0, 1])$ .

2. Let  $f \in C(\mathbb{B}(0, 1), \mathbb{R}^n)$  be such that

$$|f(x)| \leq 1 \text{ for } |x| = 1.$$

Show that  $f$  has a fixed point in  $\mathbb{B}(0, 1)$ .

3. Let  $\Omega \subset \mathbb{R}^n$  be a bounded, open set with smooth boundary. Prove that the equation

$$\begin{cases} -\Delta u = e^{-u} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

possesses a solution.

4. (*Kolmogoroff*) Show that a subset  $K \subset L_p(\mathbb{R}^n)$  ( $1 \leq p < \infty$ ) is compact iff

(i)  $K$  is closed and bounded.

(ii)  $\int_{|x| \geq N} |f(x)|^p dx \rightarrow 0$ ,  $N \rightarrow \infty$ , uniformly in  $f \in K$ .

(iii)  $\int_{\mathbb{R}^n} |f(x+h) - f(x)|^p dx \rightarrow 0$ ,  $|h| \rightarrow 0$ , uniformly in  $f \in K$ .

[Hint: Use the density of test functions in  $L_p(\mathbb{R}^n)$ , the strong continuity of the translation semigroup on  $L_p(\mathbb{R}^n)$  and Arzela-Ascoli.]

5. Let  $1 \leq p < \infty$  and prove that

$$\left( \int_{\mathbb{R}^n} |u(x+h) - u(x)|^p dx \right)^{1/p} \leq |h| \|u\|_{1,p}$$

for  $u \in W_p^1(\mathbb{R}^n)$ . Use this estimate and Kolmogoroff's characterization of compactness to show that  $W_p^1(\Omega) \hookrightarrow L_p(\Omega)$  for  $\Omega \subset \mathbb{R}^n$  open and bounded.

Homework due by Friday, May 13 2005.