

GENERALIZED SNELL'S LAW FOR WEIGHTED MINIMAL SURFACE IN HETEROGENEOUS MEDIA

ZHILIN LI AND XIAOBIAO LIN*, MONICA TORRES†, AND HONGKAI ZHAO‡

Abstract. The weighted minimal surface problem in piecewise smooth media is studied in this paper. The solution to the weighted minimal surface problem is continuous but the derivatives have a jump across the interface where the medium property is discontinuous. The jump condition of the derivatives derived in this paper generalized the Snell's law in geometric optics to weighted minimal surfaces of co-dimension one in any number of dimensions. A numerical method based on the gradient flow and the maximum principal preserving immersed interface method is developed to solve this nonlinear elliptic problem with jump conditions. Numerical computations are presented to verify both the analysis and the numerical algorithm.

1. Introduction. The minimal surface problem, that is, the problem of finding the surface of the least area among all surfaces having fixed boundary data, has been extensively studied. Many phenomena that occur in nature relate to this problem which has been a motivation for developing new mathematical theories and techniques to solve the problem analytically and numerically. A recent workshop on minimal surfaces presented the latest research on minimal surface applications in chemistry and biology [5]. Minimal surfaces were shown to be important in various chemical micro-structures and their corresponding phase transitions [5]. Computer graphics and image analysis use minimal surfaces frequently for boundary detection, and to construct surfaces that are visually appealing [2], [13]. Soap films and other membranes passing through a fixed boundary provide mechanical examples of minimal surfaces [12]. A related concept is the idea of *capillary surfaces*, which result from surface tension in liquids. These surfaces are closely related to minimal surfaces [6]. For a precise mathematical description of the minimal surface problem we refer, for example, to the classical treatises [7] and [14].

The minimal surface problem can be described in two different ways, using the parametric or the non-parametric formulation. In the non-parametric setting the candidate surfaces are graphs of functions, while in the parametric setting the surfaces are treated as boundaries of sets [7]. The former is usually seen in more physically-based treatments of the problem, whereas the later provides an excellent framework for the mathematical analysis of minimal surfaces.

When the medium is homogeneous, the energy density at each point is constant, and therefore the surface energy is equivalent to the surface area. This is the standard minimal surface problem. In this paper, we consider the weighted minimal surface problem in a heterogeneous medium in which the energy density is piecewise smooth.

*Center for Research in Scientific Computation & Department of Mathematics, North Carolina State University, Raleigh, NC 27695 (zhilin@math.ncsu.edu, xblin@math.ncsu.edu).

†Department of Mathematics, Northwestern University, Evanston, IL 60208. (torres@math.northwestern.edu).

‡Department of Mathematics, University of California, Irvine, 92697 (zhao@math.uci.edu).

For example this is the case for capillary interfaces in porous media or composite materials. In particular we derive a jump condition for the weighted minimal surface at the interface between two different media. The jump condition can be regarded as a generalized Snell's law which describes the "refraction" of minimal surfaces instead of light rays in geometric optics.

For simplicity, we restrict our discussion to a two dimensional surface in three dimensions. Our results extend to minimal surfaces of co-dimension one in any number of dimensions. Below we present the setup of the problem: given a two dimensional domain $\Omega \subset R^2$, we want to find a function $u(x, y)$ which minimizes the weighted surface area of its graph with a prescribed Dirichlet boundary condition, i.e.,

$$\begin{aligned} \min_u E(u) &= \iint_{\Omega} c(x, y, u) \sqrt{1 + |\nabla u(x, y)|^2} dx dy, \\ u(x, y)|_{\partial\Omega} &= u_0(x, y), \end{aligned} \tag{1.1}$$

where $c(x, y, u)$ is a piecewise smooth function which has a finite jump across a surface defined by

$$S = \{ (x, y, z), \varphi(x, y, z) = 0 \}, \tag{1.2}$$

in three dimensional space, see Fig. 2.1 for an illustration. We assume that the surface is smooth and $\varphi(x, y, z)$ has up to second order continuous partial derivatives. In this paper we consider the case $c > 0$. Mathematically, the case when $c = 0$ has been studied in [15] using standard arguments of the calculus of variations. In [15], the minimal surfaces are treated as the boundary of sets of the finite perimeter (not graphs of functions) and the case with weight zero inside the medium and one outside is studied. In particular, minimal surfaces that always stay at a bounded distance (universal) from a given plane are constructed. For this degenerate case it can be proven ([15]) that the minimal surfaces enter the medium inside *orthogonally* which means, roughly speaking, that the intersection of the minimal surface and the boundary that separates two mediums locally looks like two perpendicular hyperplanes. Actually the orthogonality condition for the degenerate case $c = 0$ can be formally derived through taking the limit of our generalized Snell's law in this paper.

In practice the computation of a weighted minimal surface in a piecewise smooth medium is quite challenging due to the nonlinearity of the Euler-Lagrange equation and the discontinuity of the coefficient across the interface. The intersection of the minimal surface and the interface is a free boundary problem that is coupled with the Euler-Lagrange equation through the generalized Snell's law. In this paper, we have developed an iterative numerical algorithm based on the gradient flow and the immersed interface method to compute the weighted minimal surface. In particular we represent the minimal surface as a graph function and formulate the generalized Snell's law as an interface condition for the graph function to reduce the computational cost. We linearize the equation and jump condition by substituting the previous approximation in the nonlinear terms and use the maximum principle preserving scheme for parabolic interface problem [1, 10]. A smoothing technique and a re-initialization process is used to regularize the problem numerically. Since the energy

functional is not convex, there are local minima and hence the solution depends on the initial condition $u_0(x, y)$ (see Fig. 4.4).

The paper is organized as follows. In §2, we derive the Euler-Lagrange equation and the generalized Snell's law at the interface of two mediums. In §3, we present a general computational framework and outline the algorithm that is used to compute the weighted minimal surface. Some numerical examples are presented in §4. The paper is concluded in §5.

2. Derivation of the Euler-Lagrangian equation and the jump condition.

2.1. The Euler-Lagrangian Equation. When $c \equiv 1$ in the entire domain, the minimal surface problem is commonly known as Plateau's Problem [4]. In general, there may be one, multiple, or no minimal surfaces satisfy the given boundary condition. The existence of a solution to the general case proved by Douglas (1931 [4]), and Radó (1933, [11]) could not exclude the possibility of singularities. In this paper, we only consider the case that the minimal surface is smooth except at the interface. Although our derivation of the Euler-Lagrangian equation is not new, we include it in details not only because that the weight c is non-constant in our paper, but also it is needed in the second half of this section where the jump condition across the interface S is considered.

Assume that $c(x, y, z)$ is a piecewise smooth function in \mathbb{R}^3 :

$$c(x, y, z) = \begin{cases} c_2(x, y, z), & (x, y, z) \in B, \\ c_1(x, y, z), & (x, y, z) \in \mathbb{R}^3 - B, \end{cases}$$

where $B = \{ (x, y, z) \mid \varphi(x, y, z) < 0 \}$ is an open set in \mathbb{R}^3 with the smooth boundary $S = \{ (x, y, z) \mid \varphi(x, y, z) = 0 \}$. Since S is smooth, we can extend $c_1(x, y, z)$ and $c_2(x, y, z)$ smoothly across S in a small neighborhood of S from one side to the other for theoretical convenience.

Let Ω be an open set in \mathbb{R}^2 whose boundary $\partial\Omega$ is smooth. Assume that the cylindrical region $\Omega \times \mathbb{R}$ contains B . A function $u_0(x, y)$ on $\partial\Omega$ is specified. We look for a piecewise smooth function $u(x, y)$ in Ω that satisfies $u(x, y)|_{\partial\Omega} = u_0(x, y)$ and minimizes $E(u)$ in (1.1). Assume that the graph of $u(x, y)$ intersects with the surface S at a curve Γ , and the projection of Γ to the (x, y) -plane is a smooth curve γ , which divides the domain Ω into two subsets Ω_1 and Ω_2 . Assume also that the function $u(x, y)$ is smooth in each Ω_j , $j = 1, 2$. Across γ , the function $u(x, y)$ is continuous but (u_x, u_y) usually has a finite jump.

Let $\gamma(s) = \{ (x(s), y(s)) \mid a \leq s \leq b, x(a) = x(b), y(a) = y(b) \}$, and

$$F(x, y, z, z_x, z_y) = c(x, y, z) \sqrt{1 + z_x^2 + z_y^2}. \quad (2.1)$$

With $z = u(x, y)$, the variational equation of $E(u)$ is

$$\delta E(u) = \iint_{\Omega} (F_z \delta z + F_{z_x} \delta z_x + F_{z_y} \delta z_y) dx dy = 0,$$




FIG. 2.1. A diagram of the weighted minimal surface problem. The solution $u(x, y)$ is piecewise smooth but has a finite jump in the partial derivatives across γ .

where δz is a variation of z and satisfies $\delta z|_{\partial\Omega} = 0$.

We prove the main theorem in this subsection below.

THEOREM 2.1. *If $u(x, y)$ is C^2 in $\Omega_1 \cup \Omega_2$ is a local minimum of (1.1), then it satisfies the following Euler-Lagrangian equation in the open set $\Omega_1 \cup \Omega_2$:*

$$\frac{\partial}{\partial x} \left(\frac{cu_x}{\sqrt{1+u_x^2+u_y^2}} \right) + \frac{\partial}{\partial y} \left(\frac{cu_y}{\sqrt{1+u_x^2+u_y^2}} \right) = c_z(x, y, u) \sqrt{1+u_x^2+u_y^2}. \quad (2.2)$$

Proof. Assume that the graph of $u(x, y)$ passes through Γ , i.e., $\delta z|_{\gamma} = 0$. If u is a local minimizer, we must have $\delta E = 0$ for such δz . Using the Green's formula, we have

$$\begin{aligned} \delta E &= \iint_{\Omega_1} + \iint_{\Omega_2} (F_z \delta z + F_{\delta z_x} \delta z_x + F_{\delta z_y} \delta z_y) dx dy \\ &= \iint_{\Omega_1} + \iint_{\Omega_2} (F_z - \frac{\partial}{\partial x} F_{z_x} - \frac{\partial}{\partial y} F_{z_y}) \delta z dx dy \\ &= 0. \end{aligned}$$

Since δz is arbitrary in $\Omega_1 \cup \Omega_2$, we must have

$$F_z - \frac{\partial}{\partial x} F_{z_x} - \frac{\partial}{\partial y} F_{z_y} = 0, \quad \Omega_1 \cup \Omega_2. \quad (2.3)$$

The Euler-Lagrangian equation (2.2) follows by plugging F into (2.3). ■

We will assume that $z = u(x, y)$ satisfies the Euler-Lagrange equation (2.3) in the rest of the paper.

2.2. The jump condition across the interface. Let

$$\Gamma_0 = \{(x_0(s), y_0(s), z_0(s)) \mid z_0(s) = u(x_0(s), y_0(s))\}$$

be the intersection of the minimal surface and S . Let $\Gamma = \Gamma_0 + \delta\Gamma$ be a closed curve near Γ_0 . For every small perturbation $\delta\Gamma$, we assume that there is unique solution $u + \delta u$ to the Euler-Lagrangian equation in Ω_1 and Ω_2 that passes through Γ and satisfies the boundary condition at $\partial\Omega$. In this subsection we study the variation $\delta E(u)$ with respect to $\delta\Gamma$ and derive the jump condition for the solution u across γ . The result is summarized in the following theorem.

THEOREM 2.2. *Assume that, in a neighborhood of Γ , the surface $S = \{\varphi(x, y, z) = 0\}$ can be expressed as $z = \phi(x, y)$. Let*

$$\mathbf{p}_j = \frac{(-u_x, -u_y, 1)}{\sqrt{1 + u_x^2 + u_y^2}}, \quad \mathbf{q} = \frac{(-\phi_x, -\phi_y, 1)}{\sqrt{1 + \phi_x^2 + \phi_y^2}},$$

be the unit normal directions of the surface $u(x, y)$ in Ω_j , and the unit normal direction of S respectively, all pointing to the positive direction in z direction. Then if $u(x, y)$ is the solution of the weighted minimal surface problem, the jump of the derivatives of $u(x, y)$ across the surface S must satisfy the following generalized Snell's Law:

$$c_1 \mathbf{p}_1 \cdot \mathbf{q}|_\gamma = c_2 \mathbf{p}_2 \cdot \mathbf{q}|_\gamma, \quad (2.4)$$

where γ is the intersection of the minimal surface and the interface S .

Proof. To derive the jump condition for a minimizer $u(x, y)$ across γ , we shall compute the variation of $E(u)$ due to the variation of Γ .

The variation of Γ naturally decomposes into the vertical and horizontal directions. We first assume the variation of Γ is vertical, i.e.

$$\delta\Gamma = \{(0, 0, \delta z(s))\}.$$

Let $\delta E = \delta E_1 + \delta E_2$ where δE_j is the variation in Ω_j , $j = 1, 2$,

$$\delta E_j = \iint_{\Omega_j} (F_z \delta z + F_{z_x} \delta z_x + F_{z_y} \delta z_y) dx dy.$$

Using the Euler-Lagrange equation in Ω_j and the Green's formula, we have the variational equation due to the vertical variation of Γ :

$$\delta E_j = - \oint_\gamma \delta z(s) (F_{z_x}, F_{z_y}) \cdot \mathbf{n}_j(s) ds, \quad j = 1, 2, \quad (2.5)$$

where $\mathbf{n}_j(s)$ is the inward unit normal vector of Ω_j at $\gamma(s)$.

Next, we consider the variation of E_j due to the variation in Γ on the surface of the graph of $u(x, y)$. To compute δE_1 , we will move γ into Ω_1 , and to compute δE_2 , we will move γ into Ω_2 . Each δE_j will be treated as a linear functional of $\delta\Gamma$ and will be obtained from a one-sided differentiation of E_j with respect to $\delta\Gamma$ in function spaces.




FIG. 2.2. The variation of Γ and u near the interface S .

In each Ω_j , in a small neighborhood of γ , we introduce a local coordinate system

$$x = x(s, n), \quad y = y(s, n),$$

where s is the arc length along γ , and n is the distance to $\gamma(s)$ along the inward normal direction of Ω_j at $\gamma(s)$. Assume that n is sufficiently small, then the change of coordinates $(x, y) \rightarrow (n, s)$ is nonsingular and $dxdy \approx dsdn$. In fact, $dxdy = \left| \frac{\partial(x, y)}{\partial(s, n)} \right| dsdn$ and the determinant is unity at $n = 0$. Therefore, we have

$$\left| \frac{\partial(x, y)}{\partial(s, n)} \right| = 1 + O(\delta n(s)).$$

Let $\delta\Gamma(s) = (x(s, \delta n(s)), y(s, \delta n(s), \delta z(s)))$, where $\delta n(s) \geq 0$ and $\delta z(s)$ is chosen so that $\Gamma + \delta\Gamma$ is on the graph of $u(x, y)$. Denote the region bounded by γ and $\gamma + \delta\gamma$ as $\delta\Omega_j$. We perform the following derivation

$$\begin{aligned} \delta E_j &= - \int \int_{\delta\Omega_j} F \, dxdy \\ &= - \int \int_{\delta\Omega_j} F \left| \frac{\partial(x, y)}{\partial(x, n)} \right| \, dsdn \\ &= - \oint_{\gamma} \int_0^{\delta n(s)} (1 + O(\delta n(s))) F \, dn ds \\ &= - \oint_{\gamma} F(s) \delta n(s) \, ds + O((\delta n(s))^2). \end{aligned}$$

Dropping the higher order terms, we have

$$\delta E_j = - \oint_{\gamma} F(s) \delta n(s) \, ds. \quad (2.6)$$

Note that

$$\delta E_j = \frac{\partial E_j}{\partial n} \delta n + \frac{\partial E_j}{\partial z} \delta z. \quad (2.7)$$

Here $\frac{\partial E_j}{\partial n}$ and $\frac{\partial E_j}{\partial z}$ are linear functionals on δn and δz .

From (2.5), we have

$$\frac{\partial E_j}{\partial z} \delta z = - \oint_{\gamma} \left((F_{z_x}, F_{z_y}) \cdot \mathbf{n} \right) \delta z(s) ds.$$

Combining this with (2.6) and (2.7), we have

$$\frac{\partial E_j}{\partial n} \delta n = - \oint_{\gamma} F(s) \delta n(s) ds + \oint_{\gamma} \left((F_{z_x}, F_{z_y}) \cdot \mathbf{n} \right) \delta z(s) ds.$$

Since we assume that Γ is on the surface of $u(x, y)$, see Fig. 2.2(a), there is a relation between the vertical and horizontal variation:

$$\delta z(s) = \frac{\partial u}{\partial n} \delta n(s).$$

Thus we have found the variational equation due to the horizontal variation of Γ :

$$\frac{\partial E_j}{\partial n} \delta n = \oint_{\gamma} \left(-F(s) + (F_{z_x}, F_{z_y}) \cdot \mathbf{n} \frac{\partial u}{\partial n} \right) \delta n(s) ds. \quad (2.9)$$

To keep $\Gamma + \delta\Gamma$ on the surface S where c has a jump discontinuity, we assume that $\delta z(s) = \frac{\partial \phi}{\partial n} \delta n(s)$ in (2.5). So the total change in z direction due to the perturbation $\delta n(s)$ is $(\frac{\partial \phi}{\partial n} - \frac{\partial u}{\partial n}) \delta n(s)$, see Fig. 2.2(b). Thus the total variation for Γ on S is

$$\delta E_j = \oint_{\gamma} \left(-F(s) + (F_{z_x}, F_{z_y}) \cdot \mathbf{n} \left(\frac{\partial u}{\partial n} - \frac{\partial \phi}{\partial n} \right) \right) \delta n(s) ds, \quad j = 1, 2.$$

Define

$$\begin{aligned} \tilde{F}(s, n, z, z_s, z_n) &:= F(x, y, z, z_x, z_y) \\ &= F \left(x(s, n), y(s, n), z(s, n), \frac{\partial s}{\partial x} z_s + \frac{\partial n}{\partial x} z_n, \frac{\partial s}{\partial y} z_s + \frac{\partial n}{\partial y} z_n \right). \end{aligned}$$

Note that $\mathbf{n} = (\frac{\partial n}{\partial x}, \frac{\partial n}{\partial y})$. Thus we have

$$\tilde{F}_{z_n} = F_{z_x} \frac{\partial n}{\partial x} + F_{z_y} \frac{\partial n}{\partial y} = (F_{z_x}, F_{z_y}) \cdot \mathbf{n}. \quad (2.11)$$

Denoting \tilde{F} by F if no confusion should arise, we have:

$$\delta E_j = - \oint_{\gamma} \left(F(s) + F_{z_n} \left(\frac{\partial \phi}{\partial n} - \frac{\partial u}{\partial n} \right) \right) \delta n(s) ds, \quad j = 1, 2. \quad (2.12)$$

More generally, let $\delta n(s)$ be an arbitrarily function of s , not necessarily nonnegative as in the previous paragraphs. Since in Ω_1 and Ω_2 , the inward normal directions are opposite to each other, if $\delta n(s) = \psi(s)$ in Ω_1 , then $\delta n(s) = -\psi(s)$ in Ω_2 where $\psi(s)$ is an arbitrarily smooth function of s . The total variation in $\Omega_1 \cup \Omega_2$ must add up to zero, therefore, we have the jump condition across the surface S :

$$\left(F(s) + F_{z_n} \left(\frac{\partial \phi}{\partial n} - \frac{\partial u}{\partial n} \right) \right) \Big|_{j=1} = \left(F(s) + F_{z_n} \left(\frac{\partial \phi}{\partial n} - \frac{\partial u}{\partial n} \right) \right) \Big|_{j=2}. \quad (2.13)$$

We now apply (2.13) to the minimal surface problem with F given in (2.1). Using (2.11), we have

$$\begin{aligned} F_{z_n} &= \left(\frac{cu_x}{\sqrt{1+u_x^2+u_y^2}}, \frac{cu_y}{\sqrt{1+u_x^2+u_y^2}} \right) \cdot \mathbf{n} \\ &= \frac{cu_n}{\sqrt{1+u_x^2+u_y^2}}. \end{aligned} \quad (2.14)$$

Note that along the line Γ where $u(s, n)$ and $\phi(s, n)$ meet, $u(s, 0) = \phi(s, 0)$, $u_s = \phi_s$ and $1 + u_x^2 + u_y^2 = 1 + u_s^2 + u_n^2$. From (2.14), on the line Γ , for $j = 1, 2$, we have,

$$\begin{aligned} F(s) + F_{z_n} \left(\frac{\partial \phi}{\partial n} - \frac{\partial u}{\partial n} \right) &= \frac{c(x, y, u)(1 + u_s u_s + u_n \phi_n)}{\sqrt{1 + u_s^2 + u_n^2}} \\ &= \frac{c(x, y, u)(1 + u_s \phi_s + u_n \phi_n)}{\sqrt{1 + u_s^2 + u_n^2}}. \end{aligned} \quad (2.15)$$

If we multiply both sides of (2.13) by $1/\sqrt{1 + \phi_s^2 + \phi_n^2}$, which is continuous across the interface, and use $u_x \phi_x + u_y \phi_y = u_s \phi_s + u_n \phi_n$, we have finally arrived at (2.4). \blacksquare

Remarks

1. If the interface S is expressed in terms of the zero level surface $\varphi(x, y, z) = 0$, then the jump condition is equivalent to

$$\left[\frac{c(u_x \varphi_x + u_y \varphi_y)}{|\nabla \varphi| \sqrt{1 + |\nabla u|^2}} \right] = \left[\frac{c \varphi_z}{|\nabla \varphi| \sqrt{1 + |\nabla u|^2}} \right], \quad (2.16)$$

where $(\varphi_x, \varphi_y, \varphi_z)/|\nabla \varphi|$ is the unit normal direction of the surface S .

2. The jump condition (2.4) is a local condition and can be written in a coordinate free form. Define the dihedral angle θ between two surfaces to be the angle between their normals. Then the generalized Snell's law (2.4) can be expressed as

$$c_1 \cos(\theta_1) = c_2 \cos(\theta_2), \quad (2.17)$$

where θ_j is the dihedral angle between the interface S and the surface $u_j(x, y)$. Our derivations and the generalized Snell's law extend to weighted minimal surfaces of co-dimension one in any number of dimensions.

3. A numerical method. Numerical computation of weighted minimal surfaces in heterogeneous media in three dimensions is not easy. First of all the Euler-Lagrange equation is nonlinear. Moreover, we need to incorporate the generalized Snell's law across a free boundary, the intersection of the minimal surface and the interface

between two different media. The main difficulty is that tracking the free boundary, enforcing the jump condition across the free boundary, and solving the nonlinear partial differential equation in separated regions have to be handled simultaneously and consistently.

In this paper, we propose an iterative approach based on the gradient flow. We represent the minimal surface as a graph and reformulate the the generalized Snell's law as a two dimensional interface condition for the graph function so that we only need to solve the Euler Lagrange equation in *two space dimensions*, and takes advantage of the *maximum principle preserving scheme* for linear elliptic interface problems [10].

In this section, we assume that c is a *piecewise constant* with a finite jump across the interface S . We would like to compute the solution to the Euler-Lagrange equation

$$\frac{\partial}{\partial x} \left(\frac{c u_x}{\sqrt{1 + |\nabla u|^2}} \right) + \frac{\partial}{\partial y} \left(\frac{c u_y}{\sqrt{1 + |\nabla u|^2}} \right) = 0, \quad (x, y) \in \Omega_1 \cup \Omega_2 - \gamma \quad (3.1)$$

with the jump condition (2.16) directly. Due to the difficulties mentioned earlier, a direct solver to the elliptic non-linear problem is very challenging. Moreover, the solution may not be unique, and may be very sensitive to the boundary conditions and the geometric configuration of the interface, see Section 4. To prevent the approximate solution from jumping from one solution to another, we solve the following gradient flow, a parabolic equation, to steady state

$$\begin{aligned} u_t &= \frac{\partial}{\partial x} \left(\frac{c u_x}{\sqrt{1 + |\nabla u|^2}} \right) + \frac{\partial}{\partial y} \left(\frac{c u_y}{\sqrt{1 + |\nabla u|^2}} \right) \\ u(x, y, t)|_{\partial\Omega} &= u_0(x, y), \end{aligned} \quad (3.2)$$

where $u_0(x, y)$ is the boundary condition. An initial guess that satisfies the boundary condition has to be chosen and the time step can be adjusted depending on the geometry of the problem.

3.1. The outline of the algorithm. We express the boundary S between the two different mediums as the zero level set of a function $\varphi(x, y, z)$:

$$\varphi(x, y, z) \begin{cases} < 0, & \text{if } (x, y, z) \in S, \\ = 0, & \text{if } (x, y) \text{ is on } S, \\ > 0, & \text{if } (x, y) \text{ is outside of } S. \end{cases} \quad (3.3)$$

The actual computation is done on a two dimensional domain $[a, b] \times [c, d]$. We use a uniform Cartesian grid

$$x_i = a + i h_x, \quad i = 0, 1, \dots, M; \quad y_j = c + j h_y, \quad j = 0, 1, \dots, N, \quad (3.4)$$

where $h_x = \frac{b-a}{m}$ and $h_y = \frac{d-c}{n}$. We need to carefully choose an initial guess $u^0(x, y)$ that satisfies the boundary condition in order to get the intended solution. We use the substitution method, see [8] for example, to find an approximate solution to the weighted minimal surface problem. The main procedure from $u^k(\mathbf{x})$ to get $u^{k+1}(\mathbf{x})$ is outlined below:

- **Step 1:** Regularize the numerical solution by the following modified Helmholtz equation

$$u^* - \Delta t_1 \Delta u^* = u^k \quad (3.5)$$

where Δt_1 is a regularization factor and is of order h^2 , and the Laplacian operator is discretized in the standard way:

$$\Delta_h u_{ij}^* = \frac{u_{i-1,j}^* - 2u_{ij}^* + u_{i+1,j}^*}{(h_x)^2} + \frac{u_{i,j-1}^* - 2u_{ij}^* + u_{i,j+1}^*}{(h_y)^2}. \quad (3.6)$$

The purpose of this step is to smooth the intermediate approximation $u^k(x, y)$ without changing it too much. This step is introduced because the intermediate solution to the interface problem usually has non-smooth second order derivatives, and this may affect the convergence of our method. We use the fast solver from the Fishpack which is available through the Netlib to solve the above equation.

- **Step 2:** Set $\varphi_2^k(x_i, y_j) = \varphi(x_i, y_j, u_{ij}^*)$. If $\varphi_2^k(x, y) = 0$ is non-empty, that is

$$\max_{ij} \{\varphi_2^k(x_i, y_j)\} \min_{ij} \{\varphi_2^k(x_i, y_j)\} < 0,$$

we use a re-initialization scheme so that $\varphi_2^k(x, y)$ is a good approximation to the signed distance function $\varphi_2^k(x, y) = 0$, and has up to second order partial derivatives in the neighborhood of its zero level set. This is a standard technique in the level set method. The re-initialization is essential to the algorithm because $\varphi(x, y, u^*(x, y))$ is usually a non-differentiable function across the curve $\varphi(x, y, u^*(x, y)) = 0$.

- **Step 3:** Update u^{k+1} from

$$u^{k+1} = u^* + \Delta t_2 \left\{ \frac{\partial}{\partial x} \left(\frac{c u_x^{k+1}}{\sqrt{1 + |\nabla u^*|^2}} \right) + \frac{\partial}{\partial y} \left(\frac{c u_y^{k+1}}{\sqrt{1 + |\nabla u^*|^2}} \right) \right\}$$

$$u^{k+1}(x, y) \Big|_{\partial\Omega} = u_0(x, y) \quad (3.7)$$

where Δt_2 is the time step. If the zero level set $\varphi_2^k(x, y) = 0$ is non-empty, we reformulate the jump condition (2.16) across $\varphi_2^k(x, y) = 0$ as

$$[u^{k+1}] = 0,$$

$$\left[\frac{c u_n^{k+1}}{\sqrt{1 + |\nabla u^*|^2}} \right] = \left[\frac{c}{\sqrt{1 + |\nabla u^*|^2}} \frac{\varphi_z}{|\nabla \varphi|} \right]$$

$$+ \left[\frac{c}{\sqrt{1 + |\nabla u^*|^2}} \left((n_x^k - \frac{\varphi_x}{|\nabla \varphi|}) u_x^* + (n_y^k - \frac{\varphi_y}{|\nabla \varphi|}) u_y^* \right) \right], \quad (3.8)$$

where $\mathbf{n}^k = (n_x^k, n_y^k)$ is the unit normal direction of the plane curve $\varphi_2^k(x, y) = \varphi(x, y, u^*(x, y)) = 0$ pointing outward. The gradient $\nabla\varphi$ is computed at $(x, y, u^*(x, y))$. It is obvious that if u^k converges, the modified jump condition (3.8) converges to the true jump condition (2.16). Since we are only interested in the steady state solution of (3.2), the accuracy in time variable is unimportant. We use the maximum principal preserving scheme to discretize the elliptic part, that is the right hand side of equation (3.2), see [10, 1]. The discrete implicit finite difference equation is solved using the multigrid solver [3].

Note that we have re-written the jump condition to fit perfectly into our numerical scheme for elliptic interface problems described in [10]. The crucial part in the algorithm is to compute the gradient ∇u^* at staggered grid points $(x_i \pm h_x/2, y_j)$ and $(x_i, y_j \pm h_y/2)$, and certain points on the interface $\varphi_2^k(x, y) = 0$. The detailed description can be found in [8] and is not going to be repeated here. The main methodology is the least square interpolation techniques, see [8, 9].

4. Numerical examples. We first present an example in which the interface is a cylinder. In this case, we have a fixed projected interface in the x - y plane and a homogeneous jump condition across the interface because $\varphi_z \equiv 0$, $n_x^k = \varphi_x/|\nabla\varphi|$, and $n_y^k = \varphi_y/|\nabla\varphi|$. There is a unique solution and the substitution method converges, see [8] for the details. We omit the step 1 and take $\Delta t_2 = \infty$ meaning that we solve the elliptic equation in (3.7) directly. Given an initial guess $u^0(x, y)$, the substitution method simply is

$$\left(\frac{\partial}{\partial x} \frac{c u_x^{k+1}}{\sqrt{1 + |\nabla u^k|^2}} \right) + \frac{\partial}{\partial y} \left(\frac{c u_y^{k+1}}{\sqrt{1 + |\nabla u^k|^2}} \right) = 0, \quad u^{k+1}(x, y) \Big|_{\partial\Omega} = u_0(x, y)$$

$$[u^{k+1}] = 0, \quad \left[\frac{c u_n^{k+1}}{\sqrt{1 + |\nabla u^k|^2}} \right] = 0$$

In Fig. 4.1 (a)-(b), we show the mesh and contour plots of the minimal surface with $c^- = 0.01$ and $c^+ = 1$ given the initial guess

$$u^0(x, y) = -\frac{1}{2} + \frac{x+1}{2}, \quad \Omega = [-1, 1] \times [-1, 1]. \quad (4.1)$$

The Dirichlet boundary condition is $u(x, y)|_{\partial\Omega} = u^0(x, y)|_{\partial\Omega}$. The interface is the cylinder defined by $\varphi(x, y, z) = \frac{x^2}{0.6^2} + \frac{y^2}{0.5^2} - 1$. If the two mediums are the same, that is $c^+ = c^-$, then $u^0(x, y)$ is the unique solution. This has been confirmed by our code to check the consistency. When c^+ is bigger than c^- which is the case in Fig. 4.1 (a)-(b), the minimal surface would try to avoid to pass outside the cylinder as much as possible to decrease the contribution from outside. The large area inside the cylinder was balanced by the smaller weight c^- .

In Fig. 4.1 (c)-(d), we show the results of an opposite case with $c^- = 1$ and $c^+ = 0.01$. The cylinder is more complicated and is given by $r = 0.5 + 0.2 \sin(4\theta)$, $0 \leq \theta \leq 2\pi$ in polar coordinates, where $r = \sqrt{x^2 + y^2}$. Now the weighted minimal

surface tries to avoid inside as much as possible that is the case in Fig. 4.1 (c). In Fig. 4.1 (b) and (d), the dotted line is the intersection of the cylinder and the minimal surface. The number of iterations of the substitution method is typically between $10 \sim 20$ iterations with the tolerance 10^{-6} .




FIG. 4.1. Computed results with different coefficients c and different cylinders. (a)-(b): Mesh and contour plots of the computed minimal surface $z = u(x, y)$ with $c^- = 0.01$, $c^+ = 1$. The interface that separates the two mediums is the elliptic cylinder $\frac{x^2}{0.6^2} + \frac{y^2}{0.5^2} = 1$. In (b), the dotted line is the intersection of $\varphi(x, y, z) = 0$ and $\varphi(x, y, u(x, y)) = 0$. (c)-(d): Mesh and contour plots of the computed minimal surface $z = u(x, y)$ with $c^- = 1$, $c^+ = 0.01$. The interface that separates the two mediums is the cylinder $r = 0.5 + 0.2 \sin(4\theta)$ in polar coordinates.

Now we present a numerical example in which the interface is a closed surface in three dimensional space. We take the interface that separates the two mediums as a sphere $x^2 + y^2 + z^2 = 0.75^2$. This is a much tougher example than the cylinder case because the intersection of the minimal surface and the interface becomes a free boundary which changes with the solution in each iteration, so does the the projected curve $\varphi(x, y, u(x, y)) = 0$. The final solution depends on both the boundary condition and the initial guess. The weighted minimal surface may or may not pass through or touch the interface that separates the two media. In order to avoid rapid changes in the solution, we choose $\Delta t_1 = h^2/4$ and $\Delta t_2 = h^2/4$.

In Fig. 4.2 (a) and (b), we show the computed weighted minimal surface with $c^- = 2$ and $c^+ = 1$ in (a), and $c^- = 1$ and $c^+ = 5$ in (b). The Dirichlet boundary condition is taken from the function $u(x, y) = -0.4 + \frac{x+1}{2}$. The difference in c is small and the numerical method can always give convergent results independent of the initial guess.




FIG. 4.2. Computed results with different coefficients c with a sphere that separates the two mediums. The numerical results are independent of the initial guess. (a) $c^- = 2$, $c^+ = 1$. (b) $c^- = 1$, $c^+ = 5$.

As we increase the weight c^- using the same boundary condition as in Fig. 4.2 (a), there is more contribution to the energy from inside of the medium. As a result, the minimal surface will move up within the inside medium. If c^- is large enough, then the minimal surface will move up further until it is eventually barely touches the interface. However, once an intermediate solution does not touch the interface, it will not feel the difference of the mediums and the next iteration will move it towards the solution in the homogeneous medium which is a plane in our example. Because the plane passes the medium inside, it will move up again. For this case, the weighted minimal surface barely touches the medium inside, see Fig. 4.3 (b). To prevent the intermediate approximation from jumping back and forth, we solve the parabolic equation (3.7) so that the solution will not change too much at one iteration. In Fig. 4.3 (a), we show the result of an intermediate approximation to the solution while Fig. 4.3 (b) is the computed approximation which barely touches the interface. Note that, once the surface cuts through a small portion of the sphere, the curve, which is the intersection of the sphere and the surface, and its projection on the x - y plane have large curvature. This may lead to numerical difficulties in resolving the curve and result in large errors of the computed solution to the parabolic equation which is visible from Fig. 4.3 (b).

As a last test, we show the solution depends on the initial condition as well. The boundary condition is taken as a constant $u(x, y)|_{\partial\Omega} = -0.65$. The weights are $c^- = 1$ and $c^+ = 10$ so the minimal surface will try to avoid outside as much as possible. If the initial condition is taken as the same constant $u^0(x, y)|_{\partial\Omega} = -0.65$, then it stays there because it is one of the local minimum and satisfies the Euler Lagrange equation




FIG. 4.3. Computed results with $c^- = 20$ and $c^+ = 1$. The initial guess is $u^0(x, y) = -0.4 + (x + 1)/2$ (a) An intermediate solution in which the surface moves upwards to avoid large contribution from inside of the medium. (b) The accepted solution which is almost tangent of the sphere.

plus the jump condition. However, it may not be the global minimum which is given in Fig. 4.4. In order to get the global minimum, the initial guess has to be carefully chosen. In Fig. 4.4, the initial guess is chosen as a piecewise linear function

$$u^0(x, y) = \begin{cases} -0.65 + \frac{0.65(x+1)}{0.4}, & \text{if } -1 \leq x \leq -0.6, \\ 0, & \text{if } -0.6 < x \leq 0.6, \\ -0.65 + \frac{0.65(x-1)}{0.4}, & \text{if } 0.6 < x \leq 1. \end{cases} \quad (4.2)$$




FIG. 4.4. Computed results with $c^- = 1$ and $c^+ = 10$. The initial guess is given in (4.2). The initial condition is important to get the global minimum. Another obvious local minimum is the plane $u(x, y) = -0.65$.

5. Conclusion. In this paper, the weighted minimal surface problem involving two media has been studied both analytically and numerically. A generalized Snell's

law for the weighted minimal surface at the interface between two different media is derived. A numerical method is developed to compute the weighted minimal surface which satisfies a non-linear elliptic equation with jump conditions at a free boundary, the intersection of the minimal surface and the interface between two media.

Acknowledgment. We would like to thank Dr. K. Ito of North Carolina State University for some useful suggestions and discussions. The first author is partially supported by ARO grants 39676-MA and 43751-MA and NFS grants DMS-0073403 and DMS-0201094. The last author is partially supported by an ONR grant N00014-02-1-0090, a DARPA grant N00014-12-1-0603, and the Sloan Foundation Fellowship.

REFERENCES

- [1] L. M. Adams and Z. Li. The immersed interface/multigrid methods for interface problems. *SIAM J. Sci. Comput.*, 24:463–479, 2002.
- [2] P Concus, R Finn, and D.A. Editors Hoffman. *Geometric Analysis and Computer Graphics*. Mathematical Sciences Research Institute Publications, Proceedings of a workshop held May 23-25, 1988.
- [3] D. De Zeeuw. Matrix-dependent prolongations and restrictions in a blackbox multigrid solver. *J. Comput. Appl. Math.*, 33:1–27, 1990.
- [4] J. Douglas. Solution of the problem of plateau. *Trans. Amer. Math. Soc.*, 33:263–321, 1931.
- [5] E Dubois-Violette and B. Editors Pansu. *International Workshop on Geometry and Interfaces*. Aussois, France., September,17-22,1990.
- [6] R Finn. *Equilibrium Capillary Surfaces*. Springer-Verlag, 1986.
- [7] Enrico Giusti. *Minimal surfaces and functions of bounded variation*. Birkhäuser Verlag, Basel, 1984.
- [8] K. Ito and Z. Li. The immersed interface method for nonlinear interface problems with an application. submitted, 2002.
- [9] Z. Li. A fast iterative algorithm for elliptic interface problems. *SIAM J. Numer. Anal.*, 35:230–254, 1998.
- [10] Z. Li and K. Ito. Maximum principle preserving schemes for interface problems with discontinuous coefficients. *SIAM J. Sci. Comput.*, 23:1225–1242, 2001.
- [11] T. Radó. On the Problem of Plateau. *Ergebn. d. Math. u. ihrer Grenzgebiete*, Berlin: Springer-Verlag, 1933.
- [12] G Shapiro. *The mathematics of soap films: explorations with Maple*. American Mathematical Society, 2000.
- [13] G Shapiro. *Geometric Partial Differential Equations and Image Analysis*. Cambridge University Press, 2001.
- [14] Leon Simon. *Lectures on geometric measure theory*. Australian National University Centre for Mathematical Analysis, Canberra, 1983.
- [15] Monica Torres. Plane-like minimal surfaces in periodic media with exclusions. Submitted to SIAM, Journal on Math. Analysis.