

## Putnam problems A1

2006. Find the volume of the region of points  $(x, y, z)$  such that

$$(x^2 + y^2 + z^2 + 8)^2 \leq 36(x^2 + y^2).$$

2005. Show that every positive integer is a sum of one or more numbers of the form  $2^r 3^s$ , where  $r$  and  $s$  are nonnegative integers and no summand divides another. (For example,  $23 = 9 + 8 + 6$ .)

2004. Basketball star Shanille O'Keal's team statistician keeps track of the number,  $S(N)$ , of successful free throws she has made in her first  $N$  attempts of the season. Early in the season,  $S(N)$  was less than 80% of  $N$ , but by the end of the season,  $S(N)$  was more than 80% of  $N$ . Was there necessarily a moment in between when  $S(N)$  was exactly 80% of  $N$ ?

2003. Let  $n$  be a fixed positive integer. How many ways are there to write  $n$  as a sum of positive integers,  $n = a_1 + a_2 + \cdots + a_k$ , with  $k$  an arbitrary positive integer and  $a_1 \leq a_2 \leq \cdots \leq a_k \leq a_1 + 1$ ? For example, with  $n = 4$  there are four ways: 4, 2+2, 1+1+2, 1+1+1+1.

2002. Let  $k$  be a fixed positive integer. The  $n$ -th derivative of  $\frac{1}{x^k-1}$  has the form  $\frac{P_n(x)}{(x^k-1)^{n+1}}$  where  $P_n(x)$  is a polynomial. Find  $P_n(1)$ .

2001. Consider a set  $S$  and a binary operation  $*$ , i.e., for each  $a, b \in S$ ,  $a * b \in S$ . Assume  $(a * b) * a = b$  for all  $a, b \in S$ . Prove that  $a * (b * a) = b$  for all  $a, b \in S$ .

2000. Let  $A$  be a positive real number. What are the possible values of  $\sum_{j=0}^{\infty} x_j^2$ , given that  $x_0, x_1, \dots$  are positive numbers for which  $\sum_{j=0}^{\infty} x_j = A$ ?

1999. Find polynomials  $f(x), g(x)$ , and  $h(x)$ , if they exist, such that for all  $x$ ,

$$|f(x)| - |g(x)| + h(x) = \begin{cases} -1 & \text{if } x < -1 \\ 3x + 2 & \text{if } -1 \leq x \leq 0 \\ -2x + 2 & \text{if } x > 0. \end{cases}$$

1998. A right circular cone has base of radius 1 and height 3. A cube is inscribed in the cone so that one face of the cube is contained in the base of the cone. What is the side-length of the cube?

1997. A rectangle,  $HOMF$ , has sides  $HO = 11$  and  $OM = 5$ . A triangle  $ABC$  has  $H$  as the intersection of the altitudes,  $O$  the center of the circumscribed circle,  $M$  the midpoint of  $BC$ , and  $F$  the foot of the altitude from  $A$ . What is the length of  $BC$ ?

1996. Find the least number  $A$  such that for any two squares of combined area 1, a rectangle of area  $A$  exists such that the two squares can be packed in the rectangle (without interior overlap). You may assume that the sides of the squares are parallel to the sides of the rectangle.

1995. Let  $S$  be a set of real numbers which is closed under multiplication (that is, if  $a$  and  $b$  are in  $S$ , then so is  $ab$ ). Let  $T$  and  $U$  be disjoint subsets of  $S$  whose union is  $S$ . Given that the product of any *three* (not necessarily distinct) elements of  $T$  is in  $T$  and that the product of any three elements of  $U$  is in  $U$ , show that at least one of the two subsets  $T, U$  is closed under multiplication.

1994. Suppose that a sequence  $a_1, a_2, a_3, \dots$  satisfies  $0 < a_n \leq a_{2n} + a_{2n+1}$  for all  $n \geq 1$ . Prove that the series  $\sum_{n=1}^{\infty} a_n$  diverges.

1993. The horizontal line  $y = c$  intersects the curve  $y = 2x - 3x^3$  in the first quadrant as in the figure. Find  $c$  so that the areas of the two shaded regions are equal. [Figure not included. The first region is bounded by the  $y$ -axis, the line  $y = c$  and the curve; the other lies under the curve and above the line  $y = c$  between their two points of intersection.]

1992. Prove that  $f(n) = 1 - n$  is the only integer-valued function defined on the integers that satisfies the following conditions.

- (i)  $f(f(n)) = n$ , for all integers  $n$ ;
- (ii)  $f(f(n+2) + 2) = n$  for all integers  $n$ ;
- (iii)  $f(0) = 1$ .

1991. A  $2 \times 3$  rectangle has vertices as  $(0, 0)$ ,  $(2, 0)$ ,  $(0, 3)$ , and  $(2, 3)$ . It rotates  $90^\circ$  clockwise about the point  $(2, 0)$ . It then rotates  $90^\circ$  clockwise about the point  $(5, 0)$ , then  $90^\circ$  clockwise about the point  $(7, 0)$ , and finally,  $90^\circ$  clockwise about the point  $(10, 0)$ . (The side originally on the  $x$ -axis is now back on the  $x$ -axis.) Find the area of the region above the  $x$ -axis and below the curve traced out by the point whose initial position is  $(1, 1)$ .

1990. Let

$$T_0 = 2, T_1 = 3, T_2 = 6,$$

and for  $n \geq 3$ ,

$$T_n = (n + 4)T_{n-1} - 4nT_{n-2} + (4n - 8)T_{n-3}.$$

The first few terms are

$$2, 3, 6, 14, 40, 152, 784, 5168, 40576.$$

Find, with proof, a formula for  $T_n$  of the form  $T_n = A_n + B_n$ , where  $\{A_n\}$  and  $\{B_n\}$  are well-known sequences.

1989. How many primes among the positive integers, written as usual in base 10, are alternating 1's and 0's, beginning and ending with 1?

1988. Let  $R$  be the region consisting of the points  $(x, y)$  of the cartesian plane satisfying both  $|x| - |y| \leq 1$  and  $|y| \leq 1$ . Sketch the region  $R$  and find its area.

1987. Curves  $A, B, C$  and  $D$  are defined in the plane as follows:

$$\begin{aligned} A &= \left\{ (x, y) : x^2 - y^2 = \frac{x}{x^2 + y^2} \right\}, \\ B &= \left\{ (x, y) : 2xy + \frac{y}{x^2 + y^2} = 3 \right\}, \\ C &= \{ (x, y) : x^3 - 3xy^2 + 3y = 1 \}, \\ D &= \{ (x, y) : 3x^2y - 3x - y^3 = 0 \}. \end{aligned}$$

Prove that  $A \cap B = C \cap D$ .

1986. Find, with explanation, the maximum value of  $f(x) = x^3 - 3x$  on the set of all real numbers  $x$  satisfying  $x^4 + 36 \leq 13x^2$ .

1985. Determine, with proof, the number of ordered triples  $(A_1, A_2, A_3)$  of sets which have the property that

- (i)  $A_1 \cup A_2 \cup A_3 = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$ , and
- (ii)  $A_1 \cap A_2 \cap A_3 = \emptyset$ .

Express your answer in the form  $2^a 3^b 5^c 7^d$ , where  $a, b, c, d$  are nonnegative integers.

## Putnam problems B1

2006. Show that the curve  $x^3 + 3xy + y^3 = 1$  contains only one set of three distinct points,  $A$ ,  $B$ , and  $C$ , which are vertices of an equilateral triangle, and find its area.

2005. Find a nonzero polynomial  $P(x, y)$  such that  $P(\lfloor a \rfloor, \lfloor 2a \rfloor) = 0$  for all real numbers  $a$ . (Note:  $\lfloor \nu \rfloor$  is the greatest integer less than or equal to  $\nu$ .)

2004. Let  $P(x) = c_n x^n + c_{n-1} x^{n-1} + \cdots + c_0$  be a polynomial with integer coefficients. Suppose that  $r$  is a rational number such that  $P(r) = 0$ . Show that the  $n$  numbers

$$c_n r, c_n r^2 + c_{n-1} r, c_n r^3 + c_{n-1} r^2 + c_{n-2} r, \\ \dots, c_n r^n + c_{n-1} r^{n-1} + \cdots + c_1 r$$

are integers.

2003. Do there exist polynomials  $a(x), b(x), c(y), d(y)$  such that

$$1 + xy + x^2 y^2 = a(x)c(y) + b(x)d(y)$$

holds identically?

2002. Shanille O'Keal shoots free throws on a basketball court. She hits the first and misses the second, and thereafter the probability that she hits the next shot is equal to the proportion of shots she has hit so far. What is the probability she hits exactly 50 of her first 100 shots?

2001. Let  $n$  be an even positive integer. Write the numbers  $1, 2, \dots, n^2$  in the squares of an  $n \times n$  grid so that the  $k$ -th row, from left to right, is

$$(k-1)n + 1, (k-1)n + 2, \dots, (k-1)n + n.$$

Color the squares of the grid so that half of the squares in each row and in each column are red and the other half are black (a checkerboard coloring is one possibility). Prove that for each coloring, the sum of the numbers on the red squares is equal to the sum of the numbers on the black squares.

2000. Let  $a_j, b_j, c_j$  be integers for  $1 \leq j \leq N$ . Assume for each  $j$ , at least one of  $a_j, b_j, c_j$  is odd. Show that there exist integers  $r, s, t$  such that  $ra_j + sb_j + tc_j$  is odd for at least  $4N/7$  values of  $j$ ,  $1 \leq j \leq N$ .

1999. Right triangle  $ABC$  has right angle at  $C$  and  $\angle BAC = \theta$ ; the point  $D$  is chosen on  $AB$  so that  $|AC| = |AD| = 1$ ; the point  $E$  is chosen on  $BC$  so that  $\angle CDE = \theta$ . The perpendicular to  $BC$  at  $E$  meets  $AB$  at  $F$ . Evaluate  $\lim_{\theta \rightarrow 0} |EF|$ .

1998. Find the minimum value of

$$\frac{(x + 1/x)^6 - (x^6 + 1/x^6) - 2}{(x + 1/x)^3 + (x^3 + 1/x^3)}$$

for  $x > 0$ .

1997. Let  $\{x\}$  denote the distance between the real number  $x$  and the nearest integer. For each positive integer  $n$ , evaluate

$$F_n = \sum_{m=1}^{6n-1} \min\left(\left\{\frac{m}{6n}\right\}, \left\{\frac{m}{3n}\right\}\right).$$

(Here  $\min(a, b)$  denotes the minimum of  $a$  and  $b$ .)

1996. Define a **selfish** set to be a set which has its own cardinality (number of elements) as an element. Find, with proof, the number of subsets of  $\{1, 2, \dots, n\}$  which are *minimal* selfish sets, that is, selfish sets none of whose proper subsets is selfish.

1995. For a partition  $\pi$  of  $\{1, 2, 3, 4, 5, 6, 7, 8, 9\}$ , let  $\pi(x)$  be the number of elements in the part containing  $x$ . Prove that for any two partitions  $\pi$  and  $\pi'$ , there are two distinct numbers  $x$  and  $y$  in  $\{1, 2, 3, 4, 5, 6, 7, 8, 9\}$  such that  $\pi(x) = \pi(y)$  and  $\pi'(x) = \pi'(y)$ . [A *partition* of a set  $S$  is a collection of disjoint subsets (parts) whose union is  $S$ .]

1994. Find all positive integers  $n$  that are within 250 of exactly 15 perfect squares.

1993. Find the smallest positive integer  $n$  such that for every integer  $m$  with  $0 < m < 1993$ , there exists an integer  $k$  for which

$$\frac{m}{1993} < \frac{k}{n} < \frac{m+1}{1994}.$$

1992. Let  $S$  be a set of  $n$  distinct real numbers. Let  $A_S$  be the set of numbers that occur as averages of two distinct elements of  $S$ . For a given  $n \geq 2$ , what is the smallest possible number of elements in  $A_S$ ?

1991. For each integer  $n \geq 0$ , let  $S(n) = n - m^2$ , where  $m$  is the greatest integer with  $m^2 \leq n$ . Define a sequence  $(a_k)_{k=0}^{\infty}$  by  $a_0 = A$  and  $a_{k+1} = a_k + S(a_k)$  for  $k \geq 0$ . For what positive integers  $A$  is this sequence eventually constant?

1990. Find all real-valued continuously differentiable functions  $f$  on the real line such that for all  $x$ ,

$$(f(x))^2 = \int_0^x [(f(t))^2 + (f'(t))^2] dt + 1990.$$

1989. A dart, thrown at random, hits a square target. Assuming that any two parts of the target of equal area are equally likely to be hit, find the probability that the point hit is nearer to the center than to any edge. Express your answer in the form  $\frac{a\sqrt{b} + c}{d}$ , where  $a, b, c, d$  are integers.

1988. A *composite* (positive integer) is a product  $ab$  with  $a$  and  $b$  not necessarily distinct integers in  $\{2, 3, 4, \dots\}$ . Show that every composite is expressible as  $xy + xz + yz + 1$ , with  $x, y, z$  positive integers.

1987. Evaluate

$$\int_2^4 \frac{\sqrt{\ln(9-x)} dx}{\sqrt{\ln(9-x)} + \sqrt{\ln(x+3)}}.$$

1986. Inscribe a rectangle of base  $b$  and height  $h$  in a circle of radius one, and inscribe an isosceles triangle in the region of the circle cut off by one base of the rectangle (with that side as the base of the triangle). For what value of  $h$  do the rectangle and triangle have the same area?

1985. Let  $k$  be the smallest positive integer for which there exist distinct integers  $m_1, m_2, m_3, m_4, m_5$  such that the polynomial

$$p(x) = (x - m_1)(x - m_2)(x - m_3)(x - m_4)(x - m_5)$$

has exactly  $k$  nonzero coefficients. Find, with proof, a set of integers  $m_1, m_2, m_3, m_4, m_5$  for which this minimum  $k$  is achieved.